

**NEET** | **JEE**  
CLASS XI-XII ESSENTIALS

**ACE** YOUR **WAY** **CBSE**  
Class XI-XII

# CHEMISTRY

India's #1  
CHEMISTRY MONTHLY FOR  
JEE (Main & Advanced) & NEET

 **today**

**CHEMISTRY  
MUSING**

**CONCEPT  
MAP**

**EXAMINER'S  
MIND**

**MONTHLY  
PRACTICE  
PROBLEMS  
(XI & XII)**

**CONCEPT  
BOOSTER**

**mtg**

Trust of more than  
1 Crore Readers  
Since 1982



2017100009290

**ADVANCED  
CHEMISTRY  
BLOC**



संस्कार से सफलता तक

Together,  
we will make  
a difference



Sitting: Govind Maheshwari (Director), Rajesh Maheshwari (Director)  
Standing: Naveen Maheshwari (Director), Brajesh Maheshwari (Director)

## What Makes **ALLEN** INDIA's Most Trusted Career Coaching Institute...

- ✓ 29 Years of Leadership
- ✓ Trust of 6.50 Lac+ Students & their Parents
- ✓ 4600+ Team members including 350+ IITians and 70+ Doctors
- ✓ Unmatched Education System with Indian Values
- ✓ With Maximum Talented Students joining ALLEN, it becomes a National Level Competition provider
- ✓ Only Institute giving Quality Results in both Pre-Medical & Pre-Engineering year after year
- ✓ Historical Result in AIIMS 2017- AIR 1 to 10 (All Top 10 AIR) Secured by Students of ALLEN
- ✓ All India Ranks 1, 2, 3 in both IIT-JEE (Adv.) & NEET 2016 from Classroom, Breaking all Records
- ✓ Limca Book of Records (2014) recognized ALLEN as Largest Educational (coaching) Institute of India by student strength (66,504) at a single location KOTA

**AIIMS 2017**

**FIRST TIME  
IN THE  
HISTORY**

All Top 10 AIR Secured  
by Students of ALLEN  
34 in Top 50 AIR



**NISHITA PUROHIT**  
Classroom

**NEET 2017**

6 in Top 10 AIR  
54 in Top 100 NEET OVERALL RANKS



**ARCHIT GUPTA**  
Classroom

**MANISH MULCHANDANI**  
Classroom

**ABHISHEK DOGRA**  
Classroom

**IIT-JEE 2017**

25 in Top 100 AIR  
130 in Top 500 AIR



**ONKAR M.D.**  
Distance

**RACHIT BANSAL**  
Classroom

**LAKSHAY SHARMA**  
Classroom



KOTA | AHMEDABAD | BENGALURU | BHILWARA | CHANDIGARH | INDORE  
JAIPUR | MUMBAI | RAJKOT | RANCHI | SURAT | VADODARA

Corporate Office : "SANKALP", CP-6, Indra Vihar, Kota (Rajasthan)-324005, India

WANT TO KNOW MORE,  
COME TO **ALLEN** AND EXPERIENCE THE LEADERSHIP

JEE (Advanced) | JEE (Main) | NEET-UG/AIIMS | KVPY, NTSE & Olympiads | Class 6<sup>th</sup> to 12<sup>th</sup>

[allen.ac.in](http://allen.ac.in) | [dlp.allen.ac.in](http://dlp.allen.ac.in) | [tab.allen.ac.in](http://tab.allen.ac.in)

[info@allen.ac.in](mailto:info@allen.ac.in)

+91-744-5156100

सत्य परेशान हो सकता है, पराजित नहीं

# To be a NEET champion, you need help from a CHAMPION



**mtg**

Skill. Passion. Hard work and determination. As a student sitting for the highly competitive NEET, you need all that. However, only a few will win, very likely with the help of a champion coach.

MTG's Champion Series for NEET is just the coach you need. It will guide you in identifying what's important for success and what's not. And then help you check your readiness with its most comprehensive

question bank. So you know your strengths and weaknesses right from the word go and course-correct accordingly. Put simply, MTG's Champion Series will help you manage your preparation effort for NEET for maximum outcome. The best part is you study at a pace you're comfortable with. Because it's all chapterwise, topicwise.



NCERT-based • Chapterwise • Topicwise • 10 years' solved previous test papers (all major medical entrance exams) • Concise summary at the start of each chapter for quick revision of key concepts • Analysis of importance of topics basis historical examination pattern • Test papers for self-assessment

Visit [www.MTG.in](http://www.MTG.in) to buy online.  
Or visit a leading bookseller near you.  
For more information, call 1800 300 23355 (toll-free) or 0124-6601200 today.  
Email [info@mtg.in](mailto:info@mtg.in)



# EXCEL ACADAMICS®

Bengaluru

A Premier Institute in KARNATAKA

Integrated PUC | PUC+CET | PUC+IIT-JEE | PUC+NEET

## NEET

**1** ALL INDIA RANK **18 44 97**

**SHIVANANJA S**  
H. T. No.: 61718963  
All India Quota Cat.

**PRAJWAL. K**  
H. T. No.: 61717312  
All India Quota Cat.

**CHETHAN. M**  
H. T. No.: 61718306  
All India Quota Cat.

**SHAMANTH M**  
H. T. No.: 903604299

## CET Ranks

**18 36 66**

**V. CHETAN**

**NAVYA N**  
H. T. No.: SH213

**SACHIN V A**  
H. T. No.: WB101

## JEE(Main) Score

**222 192 187**

**SACHIN VA**

**MADHU J N**

**SANJAY GOWDA M**

**227 181 172**

**DHEERAJ R**

**V CHETAN**

**NAVEEN RAJ**

**Helpline : 7676-91-7777, 7676-41-6666**

CET, IIT-JEE (Main & Advanced), NEET, AIIMS, JIPMER, COMED-K, BITS, VIT, MAHE

## REGISTRATION OPEN

**EXCLUSIVE  
NEET-2018**

New Batch

PCB  
Classes with Daily test

**K-CET / NEET / JEE  
CRASH COURSE  
2018**

2<sup>nd</sup> PUC/12th+2 Studying Students

**46 Days Residential  
SUMMER  
COURSE**

1 PUC/11th, +1, Studying Students

**1 Month Residential  
BRIDGE  
COURSE**

9<sup>th</sup> & 10<sup>th</sup> Studying Students (Maths & Science)

**Fully Solved  
OFF-LINE TEST  
SERIES**

CET, IIT-JEE (Main & Advanced),  
NEET - 2018 & 2019  
₹ 3,540/- ONLY

**Fully Solved  
ON-LINE TEST  
SERIES**

CET, IIT-JEE (Main & Advanced),  
NEET - 2018 & 2019  
₹ 7,080/- ONLY

**EXCEL ACADAMICS : #326, Opp. People Tree Hospital, Sheshadripuram College Road,  
Yelahanka New Town - 560064, Bangalore, KARNATAKA**

**Contact: 9535656277 / 9880286824 / 9900836461 / 9036357499**



To get free Mock CET/ NEET/JEE SMS your complete postal address to 7676917777









**Separate Deluxe Hostel for Boys and Girls**







## CBSE CHAMPION Chapterwise -Topicwise Solved Papers



CBSE CHAMPION Chapterwise -Topicwise Solved Papers Series contains topicwise questions and solutions asked over last decade in CBSE-Board examination.

Questions are supported with topicwise graphical analysis of previous years CBSE Board questions as well as comprehensive and lucid theory. The questions in each topic have been arranged in descending order as per their marking scheme. Questions from Delhi, All India, Foreign and Compartment papers are included. This ensures that all types of questions that are necessary for Board exam preparation have been covered.

Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme. CBSE sample paper and practice papers are also supplemented.

Examination papers for Class-10 and 12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email [info@mtg.in](mailto:info@mtg.in)

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!



**“**  
*We have reaped the benefits  
of starting early.  
Being the toppers,  
we recommend you  
to do the same.*  
**”**



**1**  
ALL INDIA RANK  
**KALPIT VEERWAL**  
JEE MAIN

**6**  
ALL INDIA RANK  
**SAURAV YADAV**  
JEE ADVANCED

**4**  
ALL INDIA RANK  
**SANKEERTH S**  
NEET

### Academic Benefits\*

- More than 800 Academic Hours & 500 Classes
- More than 15000 Academic Questions
- More than 100 Testing Hours

### Financial Benefits\*

- Upto Rs. 35000+ Saving on 1 Year Course Fee
- 50% Concession on Admission Form Fee
- Upto 90% Scholarship on Course Fee

\*T & C Apply

## ADMISSION ANNOUNCEMENT

Enroll Now for Academic Session 2018-19 at Coaching Fee of 2017-18  
**Classroom Contact Programs for Class V to XII**

Target: JEE (Main+Advanced) | JEE (Main) | AIIMS/ NEET | Pre-foundation

## TEST DATES

**29<sup>th</sup> October 2017**  
**12<sup>th</sup> & 26<sup>th</sup> November 2017**

### Resonance Eduventures Limited

**Registered & Corporate Office:** CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005

**Tel. No.:** 0744-3012100, 3012222, 6635555 | **CIN:** U80302RJ2007PLC024029

To know more: sms RESO at 56677 | website: [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | e-mail: [www.resonance.ac.in](http://www.resonance.ac.in)

**Toll Free: 1800 258 5555** | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)



# CHEMISTRY today

Volume 26

No. 11

November 2017

## Managing Editor

Mahabir Singh

## Editor

Anil Ahlawat

## Corporate Office:

Plot 99, Sector 44 Institutional area, Gurgaon - 122 003 (HR).

Tel : 0124-6601200 e-mail : info@mtg.in website : www.mtg.in

## Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital, New Delhi - 110029.

## Class 11

NEET   JEE Essentials	8
Examiner's Mind	22
Ace Your Way CBSE	31
MPP-7	38
Concept Map	46

## Class 12

NEET   JEE Essentials	41
Concept Map	47
Examiner's Mind	58
Ace Your Way CBSE	67
MPP-7	74

## Competition Edge

Chemistry Musing Problem Set 52	78
Nobel Prize 2017	80
Concept Booster	81
Advanced Chemistry Bloc	83
Chemistry Musing Solution Set 51	85

# CONTENTS

Subscribe online at [www.mtg.in](http://www.mtg.in)

Individual Subscription Rates				Combined Subscription Rates			
	1 yr.	2 yrs.	3 yrs.		1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775	PCM	900	1500	1900
Chemistry Today	330	600	775	PCB	900	1500	1900
Physics For You	330	600	775	PCMB	1000	1800	2300
Biology Today	330	600	775				

Send D.D/M.O in favour of MTG Learning Media (P) Ltd.

Payments should be made directly to : MTG Learning Media (P) Ltd,

Plot No. 99, Sector 44, Gurgaon - 122003 (Haryana)

We have not appointed any subscription agent.

Owned, Printed and Published by MTG Learning Media Pvt. Ltd. 406, Taj Apartment, New Delhi - 29 and printed by HT Media Ltd., B-2, Sector-63, Noida, UP-201307. Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.

Editor : Anil Ahlawat

Copyright © MTG Learning Media (P) Ltd.

All rights reserved. Reproduction in any form is prohibited.



# NEET | JEE

## ESSENTIALS

Class  
XI

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

## Unit 5

## Hydrogen | The s-Block Elements

### HYDROGEN

#### INTRODUCTION

Hydrogen is the first and lightest element in the periodic table. It was discovered by Henry Cavendish in 1766. He prepared the gas by treating iron with dil.  $\text{H}_2\text{SO}_4$ . Name 'hydrogen' was proposed by Lavoisier because it produces water on burning with oxygen (in Greek : hydra means water, gene means producing).

#### H Hydrogen

Symbol : H

Atomic Number : 1

Electronic configuration :  $1s^1$

Molecular forms :  $\text{H}_2$ ,  $\text{D}_2$ ,  $\text{T}_2$ , HD

Isotopes : Protium,  ${}^1_1\text{H}$ ; Deuterium,  ${}^2_1\text{H}$   
or D; Tritium,  ${}^3_1\text{H}$  or T

Ionic forms :  $\text{H}^-$ ,  $\text{H}^+$ ,  $\text{H}_2^+$ ,  $\text{H}_3^+$

#### Resemblance with alkali metals due to

1. Electronic configuration ( $ns^1$ ):  $1s^1$
2. Electropositive character, ( $M^+$ ):  $\text{H}^+$
3. Oxidation state (+1):  ${}^{+1}_{-1}\text{HCl}$ ,  ${}^{+1}_{-1}\text{NaCl}$

#### Resemblance with halogens due to

1. Electronic configuration : One  $e^-$  less than the nearest inert gas configuration.
2. Electronegative character : Both have tendency to accept one  $e^-$  to form anions.
3. Ionisation energy : Comparable with halogens.

#### Differences from alkali metals and halogens

1. Less electropositive than alkali metals and less electronegative than halogens.
2. Nature of oxides : Oxide of hydrogen is neutral *i.e.*  $\text{H}_2\text{O}$  while alkali metals form basic and halogens form acidic oxides.
3. Size of ions :  $\text{H}^+$  is much smaller than alkali metal ions and  $\text{H}^-$  is much larger than halide ions.


Position of Hydrogen in Periodic Table



# Last-minute check on your NEET readiness



₹ 550



MTG's NEET Explorer helps students self-assess their readiness for success in NEET. Attempting the tests put together by MTG's experienced team of editors and experts strictly on the NEET pattern and matching difficulty levels, students can easily measure their preparedness for success. Order now!



Scan now with your smartphone or tablet\*

## HIGHLIGHTS:

- 10 Model Test Papers based on latest NEET syllabus
- Last 15 years' solved test papers of AIPMT / NEET
- Includes NEET 2017 solved paper
- Detailed solutions for self-assessment and to practice time management



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: [info@mtg.in](mailto:info@mtg.in)


\*Application to read QR codes required

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!

## DIHYDROGEN


### Preparation

- Laboratory Method :** In laboratory, hydrogen is prepared by the action of dilute sulphuric acid on granular zinc in the Woulfe bottle.




- Commercial methods :**

- By electrolysis of water




- From syngas (Bosch process)




This process is called coal gasification.

The production of dihydrogen can be increased by reacting carbon monoxide of syngas mixtures with steam in the presence of iron chromate as catalyst.




This is called water-gas shift reaction.


Overall process can be represented as :



- Lane's Process**



- As a byproduct :** In the manufacture of NaOH and  $\text{Cl}_2$  by electrolysis of brine solution, hydrogen is produced as a byproduct.



### A new photocatalyst for extraction of hydrogen from seawater !


Hydrogen can be produced to power fuel cells by extracting the gas from seawater, but the electricity required to do it makes the process costly. Recently, UCF researchers has come up with a new hybrid nanomaterial that harnesses solar energy and uses it to generate hydrogen from seawater more cheaply and efficiently than current materials. This catalyst not only harvest a much broader spectrum of light than other materials, but also stand up to the harsh conditions found in seawater.

### Properties


- Physical Properties :**

- ▶ Colourless, odourless and tasteless gas.
- ▶ Non-polar in nature and slightly soluble in water.
- ▶ Liquefied under low temperature and high pressure.

- Chemical Properties :**



- Uses of hydrogen :**






## Hydrides

Hydrogen form binary hydrides with elements of *s*, *p*, (except noble gases), *d* and *f*-block.

### Ionic or Saline Hydrides

- Group-1,2 elements form ionic hydrides, *e.g.*, NaH, CaH<sub>2</sub>, CsH, SrH<sub>2</sub> etc.
- BeH<sub>2</sub>, MgH<sub>2</sub> have slightly covalent character.
- Used in synthesis of other useful hydrides.



### Metallic or Interstitial Hydrides

- d* and *f*-block elements form metallic hydrides. These are non-stoichiometric, being deficient in hydrogen, *e.g.*, LaH<sub>2.87</sub>, YbH<sub>2.55</sub>, etc.
- Metals of group-7, 8, 9 do not form hydrides and this region of periodic table is referred as hydride gap.
- Metallic hydrides can be used as hydrogen storage media.

### Covalent or Molecular Hydrides


- p*-Block elements form molecular or covalent hydrides. These are usually volatile compounds with low m.pt. and b.pt. These are of three types :
  - Electron-deficient hydrides** : Formed by group-13 elements, *e.g.* B<sub>2</sub>H<sub>6</sub>, (AlH<sub>3</sub>)<sub>*n*</sub>, etc.
  - Electron-precise hydrides** : Formed by group-14 elements, *e.g.*, CH<sub>4</sub>, SiH<sub>4</sub>, etc.
  - Electron-rich hydrides** : Formed by group-15, 16 and 17 elements, *e.g.*, NH<sub>3</sub>, H<sub>2</sub>O, HCl, etc.

## WATER

It is a crucial compound for the survival of all life forms. Human body has about 65% water and some plants have as much as 95% of water. It is a solvent of great importance.

### Properties

- Structure of water** : Bent structure with bond angle 104.5° and *sp*<sup>3</sup> hybridisation




- Physical Properties :**

- Pure water is colourless, odourless and tasteless.

- It freezes at 0 °C and boils at 100 °C.
- It has maximum density 1 g cm<sup>-3</sup> at 4 °C.


- Chemical Properties :**

- Amphoteric nature




This self-ionisation of water is also called auto-protolysis.

- Hydrolytic reaction



- As reducing agent




- Hydrate formation : Water combine with ionic salts is called water of crystallisation and such crystals are called hydrated salts.

- Coordinated water : [Ni(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup>
- Hydrogen bonded water : CuSO<sub>4</sub>·5H<sub>2</sub>O
- Interstitial water : BaCl<sub>2</sub>·2H<sub>2</sub>O

## Hard and Soft Water


- Water free from soluble salts of Ca and Mg, and which forms lather with soap readily is called soft water.
- Hard water does not form lather with soap.

## Hardness of Water and Methods for Removal



## HYDROGEN PEROXIDE

Hydrogen peroxide was discovered by the French chemist J.L. Thenard in 1818. Its molecular formula is H<sub>2</sub>O<sub>2</sub>.





## Physical Properties

- Very pale blue liquid, soluble in water, alcohol and ether.
- It boils at 152 °C and freezes at -0.89 °C.

## Strength of H<sub>2</sub>O<sub>2</sub>

- Volume strength means volume of O<sub>2</sub> liberated at N.T.P. by the decomposition of 1 mL of H<sub>2</sub>O<sub>2</sub>. 30 volume H<sub>2</sub>O<sub>2</sub> means 30 mL of O<sub>2</sub> liberated at NTP from 1 mL H<sub>2</sub>O<sub>2</sub>.  
Volume strength =  $5.6 \times N = 11.2 \times M$

## Chemical Properties

- Decomposition  
 $2\text{H}_2\text{O}_{2(l)} \longrightarrow 2\text{H}_2\text{O}_{(l)} + \text{O}_{2(g)} : \Delta H_{(g)} = -196.0 \text{ kJ}$   
This is an example of auto-oxidation and auto-reduction.
- Acidic nature  
 $2\text{NaOH} + \text{H}_2\text{O}_2 \longrightarrow \text{Na}_2\text{O}_2 + 2\text{H}_2\text{O}$
- Oxidising character
 


In acidic medium  
 $\text{H}_2\text{O}_2 + 2\text{H}^+ + 2e^- \longrightarrow 2\text{H}_2\text{O}$

In basic medium  
 $\text{H}_2\text{O}_2 + \text{OH}^- + 2e^- \longrightarrow 3\text{OH}^-$
- Reducing character
 

In acidic medium  
 $\text{H}_2\text{O}_2 \longrightarrow 2\text{H}^+ + \text{O}_2 + 2e^-$

In basic medium  
 $\text{H}_2\text{O}_2 + 2\text{OH}^- \longrightarrow 2\text{H}_2\text{O} + \text{O}_2 + 2e^-$
- Bleaching action (due to nascent O)  
 $\text{H}_2\text{O}_2 \longrightarrow \text{H}_2\text{O} + [\text{O}]$   
 Colouring matter + [O]  $\longrightarrow$  Colourless matter


- Addition reaction



## Uses

- As bleaching agent and disinfectant.
- Mixture of H<sub>2</sub>O<sub>2</sub> and hydrazine hydrate used as rocket propellant.
- For restoring the colour of old painting.
- As an antiseptic under the name perhydrol.

## Structure



## Heavy Water (D<sub>2</sub>O)

- Prepared by prolonged electrolysis of ordinary water.
- Used to prepare deuterium compounds like  
 $\text{CaC}_2 + 2\text{D}_2\text{O} \longrightarrow \text{C}_2\text{D}_2 + \text{Ca}(\text{OD})_2$   
 $\text{SO}_3 + \text{D}_2\text{O} \longrightarrow \text{D}_2\text{SO}_4$
- Used as a moderator in nuclear reactor and in reaction mechanism.

## Dihydrogen as a fuel

- Hydrogen economy is the transportation and storage of energy in the form of liquid or gaseous dihydrogen.
- Pollutants in combustion of dihydrogen will be less than petrol.

# THE s-BLOCK ELEMENTS

## INTRODUCTION

Elements of periodic table in which the last electron enters the outermost s-orbital, are called s-block elements. They are classified in two groups :

Group	1 (alkali metals)	2 (alkaline earth metals)
Electronic configuration	$ns^1$	$ns^2$

## ELECTRONIC CONFIGURATIONS OF s-BLOCK ELEMENTS


Alkali metals			Alkaline earth metals		
Elements	Atomic Number	Electronic configuration	Elements	Atomic Number	Electronic configuration
Lithium (Li)	3	$[\text{He}] 2s^1$	Beryllium (Be)	4	$[\text{He}] 2s^2$
Sodium (Na)	11	$[\text{Ne}] 3s^1$	Magnesium (Mg)	12	$[\text{Ne}] 3s^2$
Potassium (K)	19	$[\text{Ar}] 4s^1$	Calcium (Ca)	20	$[\text{Ar}] 4s^2$

Rubidium (Rb)	37	[Kr] 5s <sup>1</sup>	Strontium (Sr)	38	[Kr] 5s <sup>2</sup>
Caesium (Cs)	55	[Xe] 6s <sup>1</sup>	Barium (Ba)	56	[Xe] 6s <sup>2</sup>
Francium (Fr)	87	[Rn] 7s <sup>1</sup>	Radium (Ra)	88	[Rn] 7s <sup>2</sup>

## GENERAL PROPERTIES OF S-BLOCK ELEMENTS

Alkali metals	Alkaline earth metals
<b>Atomic/ionic radii</b>	
<ul style="list-style-type: none"> <li>Increase on moving down from Li to Cs.</li> </ul>	<ul style="list-style-type: none"> <li>Smaller than group-1 elements and increase from Be to Ra.</li> </ul>
<b>Ionization enthalpy</b>	
<ul style="list-style-type: none"> <li>Decreases down the group as the size increases.</li> </ul>	<ul style="list-style-type: none"> <li>1<sup>st</sup> I.E. of group-2 is higher than group-1 due to small size. In general, on moving down the group I.E. decreases.</li> </ul>
<b>Hydration enthalpy</b>	
<ul style="list-style-type: none"> <li>Decreases with increase in size. Li<sup>+</sup> &gt; Na<sup>+</sup> &gt; K<sup>+</sup> &gt; Rb<sup>+</sup> &gt; Cs<sup>+</sup></li> </ul>	<ul style="list-style-type: none"> <li>Decreases with increase in size. Be<sup>2+</sup> &gt; Mg<sup>2+</sup> &gt; Ca<sup>2+</sup> &gt; Sr<sup>2+</sup> &gt; Ba<sup>2+</sup></li> </ul>
<b>Physical appearance</b>	
<ul style="list-style-type: none"> <li>Silvery white due to presence of mobile electron.</li> </ul>	<ul style="list-style-type: none"> <li>Silver white lustre but harder than alkali metals.</li> </ul>
<b>Melting and boiling point</b>	
<ul style="list-style-type: none"> <li>Decreases with increase in atomic number.</li> </ul>	<ul style="list-style-type: none"> <li>Higher than group-1 and there is no regular trend in their m.pt. and b.pt.</li> </ul>

## CHEMICAL PROPERTIES OF ALKALI METALS



## TRENDS IN ALKALI METALS


- Strength of hydroxide (basicity) :** CsOH > RbOH > KOH > NaOH > LiOH

- Stability of carbonates :** Cs<sub>2</sub>CO<sub>3</sub> > Rb<sub>2</sub>CO<sub>3</sub> > K<sub>2</sub>CO<sub>3</sub> > Na<sub>2</sub>CO<sub>3</sub> > Li<sub>2</sub>CO<sub>3</sub>
- Reactivity towards H<sub>2</sub>/X :** Li > Na > K > Rb > Cs
- M.pt. and b.pt. of halides :** MF > MCl > MBr > MI

### Uses of Alkali Metals

<p><b>Li</b></p> <ul style="list-style-type: none"> <li>In alloy formation.</li> <li>Used in armour plate and aircraft parts.</li> <li>In refining of Cu and Ni to treat rheumatism.</li> </ul>	<p><b>Na</b></p> <ul style="list-style-type: none"> <li>Used for making antiknocking compounds.</li> <li>In sodium vapour lamps.</li> <li>Used in many organic compounds formation.</li> <li>As a liquid coolant in nuclear power stations.</li> </ul>
<p><b>K</b></p> <ul style="list-style-type: none"> <li>As a fertilizer.</li> <li>In biological systems.</li> <li>In manufacturing of soap.</li> <li>As an excellent adsorbent of CO<sub>2</sub>.</li> </ul>	<p><b>Cs</b></p> <ul style="list-style-type: none"> <li>In devising photoelectric cells (solar cells).</li> </ul>

## CHEMICAL PROPERTIES OF ALKALINE EARTH METALS





## TRENDS IN ALKALINE EARTH METALS

- **Basic character/solubility :**  $\text{Be}(\text{OH})_2 < \text{Mg}(\text{OH})_2 < \text{Ca}(\text{OH})_2 < \text{Sr}(\text{OH})_2 < \text{Ba}(\text{OH})_2$
- **Lattice enthalpy :**  $\text{Be}(\text{OH})_2 > \text{Mg}(\text{OH})_2 > \text{Ca}(\text{OH})_2 > \text{Sr}(\text{OH})_2 > \text{Ba}(\text{OH})_2$
- **Thermal stability :**  $\text{BeCO}_3 < \text{MgCO}_3 < \text{CaCO}_3 < \text{SrCO}_3 < \text{BaCO}_3$
- **Hydration enthalpy :**  $\text{Be}^{2+} > \text{Mg}^{2+} > \text{Ca}^{2+} > \text{Sr}^{2+} > \text{Ba}^{2+}$
- **Basicity of oxides :**

$\text{BeO} < \text{MgO} < \text{CaO} < \text{SrO} < \text{BaO}$	Amphoteric	Weakly basic	Basic
Strongly basic			
- **Solubility of halides :**  
 $\text{BeX}_2 < \text{MgX}_2 < \text{CaX}_2 < \text{SrX}_2 < \text{BaX}_2$
- **Reducing nature :**  
 $\text{Be} < \text{Mg} < \text{Ca} < \text{Sr} < \text{Ba}$

## ANOMALOUS BEHAVIOUR

Anomalous behaviours of the first element of a group is due to

- small atomic and ionic radii
- high electronegativity and ionization enthalpy
- high polarising power
- absence of *d*-electrons in its valence shell.

## Uses of Alkaline Earth Metals

**Be**

- In alloy formation.
- In windows of X-ray tubes.
- In high strength spring.

**Mg**

- Its alloy used in aircraft construction.
- In toothpaste.
- In aluminothermy.
- In Grignard reagent.

**Ca**

- In the extractions of metals.
- In casting and forging.
- In cement and mortar.

## Diagonal Relationship

Diagonal similarity is known as diagonal relationship, is due to similarity in ionic sizes and/or charge/radius ratio of the elements.

	Group 1	Group 2	Group 3	Group 4
2 <sup>nd</sup> period	Li	Be	B	C
3 <sup>rd</sup> period	Na	Mg	Al	Si

## Points of Difference

Li with other alkali metals	Be with other alkaline earth metals
<ul style="list-style-type: none"> <li>• Much harder</li> <li>• Higher m.pt. and b.pt.</li> <li>• Li forms monoxide while others form peroxides (<math>\text{M}_2\text{O}_2</math>) and superoxides (<math>\text{MO}_2</math>).</li> <li>• Ability to form nitrides.</li> <li>• <math>\text{LiOH}</math> is weak base while others are strong.</li> <li>• Ability to form hydrates <math>\text{LiCl} \cdot 2\text{H}_2\text{O}</math></li> <li>• <math>\text{LiNO}_3</math> forms the oxide on gentle heating while others form nitrites.  <math>4\text{LiNO}_3 \rightarrow 2\text{Li}_2\text{O} + 4\text{NO}_2 + \text{O}_2</math>  <math>2\text{NaNO}_3 \rightarrow 2\text{NaNO}_2 + \text{O}_2</math> </li> </ul>	<ul style="list-style-type: none"> <li>• Harder than magnesium.</li> <li>• Higher m.pt. and b.pt.</li> <li>• Does not react with cold water while other metals (except Mg) do, e.g.,  <math>\text{Ca} + 2\text{H}_2\text{O} \rightarrow \text{Ca}(\text{OH})_2 + \text{H}_2</math> </li> <li>• Be forms covalent compound while others form ionic.</li> <li>• Does not exhibit coordination number more than four.</li> <li>• Oxide and hydroxide of Be are amphoteric in nature.</li> <li>• <math>\text{Be}_2\text{C}</math> reacts with water, forming methane while others give alkyne.  <math>\text{Be}_2\text{C} + 4\text{H}_2\text{O} \rightarrow 2\text{Be}(\text{OH})_2 + \text{CH}_4</math>  Methane  <math>\text{Mg}_2\text{C}_3 + 4\text{H}_2\text{O} \rightarrow 2\text{Mg}(\text{OH})_2 + \text{C}_3\text{H}_4</math>  Propyne </li> </ul>

## Compounds of Sodium

### Sodium Carbonate (Washing soda, $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$ )

#### Preparation :

- By solvay process.  
 $2\text{NH}_3 + \text{H}_2\text{O} + \text{CO}_2 \longrightarrow (\text{NH}_4)_2\text{CO}_3$   
 $(\text{NH}_4)_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2 \longrightarrow 2\text{NH}_4\text{HCO}_3$   
 $\text{NH}_4\text{HCO}_3 + \text{NaCl} \longrightarrow \text{NH}_4\text{Cl} + \text{NaHCO}_3$   
 $2\text{NaHCO}_3 \xrightarrow{\Delta} \text{Na}_2\text{CO}_3 + \text{CO}_2 + \text{H}_2\text{O}$

#### Properties :

- White crystalline solid, exist as  $\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}$ .
- Its anhydrous form is known as soda ash.

#### Uses :

- For softening of hard water.
- In washing.
- In textile industry.
- Metal refining.
- As a laboratory reagent.

### Sodium Hydroxide (Caustic soda, $\text{NaOH}$ )

#### Preparation :

- By electrolysis of  $\text{NaCl}$  in Castner-Kellner cell.  
Cathode :  $\text{Na}^+ + e^- \xrightarrow{\text{Hg}} \text{Na-amalgam}$   
Anode :  $\text{Cl}^- \longrightarrow 1/2\text{Cl}_2 + e^-$   
 $2\text{Na-amalgam} + 2\text{H}_2\text{O} \longrightarrow 2\text{NaOH} + 2\text{Hg} + \text{H}_2$

#### Properties :

- White, translucent and deliquescent solid, soluble in water.
- Its aqueous solution is soapy to touch and corrosive.

#### Uses :

- In manufacturing of soap, paper and artificial silk.
- In petroleum refining.
- In purification of bauxite.
- In textile industries for mercerising cotton fabrics.

### Sodium Chloride ( $\text{NaCl}$ )

#### Preparation :

- Sea water is the major source. Concentrated solution of sea water is saturated with  $\text{HCl}$  gas.

#### Properties :

- White crystalline solid, soluble in water.
- Density of pure  $\text{NaCl}$  is 2.17 g/mL.
- Pure  $\text{NaCl}$  is non-hygroscopic.
- It has high m.pt. and b.pt.
- It is insoluble in alcohol.

#### Uses :

- As a table salt.
- As a freezing mixture.
- In manufacturing of  $\text{Na}_2\text{CO}_3$ ,  $\text{NaOH}$ , etc.
- In textile industries.
- Impure salt (rock salt) is used to de-ice roads.
- As a preservative of food articles like fish, meat, etc.

### Sodium Hydrogen Carbonate (Baking powder, $\text{NaHCO}_3$ )

#### Preparation :



#### Properties :

- White crystalline solid, sparingly soluble in water.
- Aqueous solution is alkaline in nature.
- Aqueous solution is weakly basic.  
 $\text{NaHCO}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NaOH} + \text{H}_2\text{CO}_3$
- The solution gives yellow colour with methyl orange but no colour with phenolphthalein.

#### Uses :

- In fire extinguishers.
- In preparation of baking powder.
- As an antiseptic and antacid.
- In manufacturing of soda.



## COMPOUNDS OF CALCIUM

### Calcium Hydroxide (Slaked Lime) ; $\text{Ca(OH)}_2$

#### Preparation :

- $\text{CaO} + \text{H}_2\text{O} \longrightarrow \text{Ca(OH)}_2$
- $\text{CaCl}_2 + 2\text{NaOH} \longrightarrow \text{Ca(OH)}_2 + 2\text{NaCl}$

#### Properties :

- White amorphous powder, sparingly soluble in water.
- Its aqueous solution is known as lime water and suspension of it in water is known as milk of lime.
- $\text{Ca(OH)}_2 + \text{CO}_2 \longrightarrow \text{CaCO}_3 + \text{H}_2\text{O}$   
(Milkiness)
- $2\text{Ca(OH)}_2 + 2\text{Cl}_2 \longrightarrow \underbrace{\text{CaCl}_2 + \text{Ca(OCl)}_2}_{\text{Bleaching Powder}} + 2\text{H}_2\text{O}$
- $\text{CaCO}_3 + \text{H}_2\text{O} + \text{CO}_2 \longrightarrow \underbrace{\text{Ca(HCO}_3)_2}_{\text{Soluble}}$

#### Uses :

- In the manufacture of bleaching powder.
- For absorbing acidic gases.
- For detection of  $\text{CO}_2$ .
- As a disinfectant.

### Calcium Oxide (Quick Lime) ; $\text{CaO}$

#### Preparation :

- $\text{CaCO}_3 \xrightarrow{\Delta} \text{CaO} + \text{CO}_2$

#### Properties :

- It is a white amorphous solid.
- On heating in oxyhydrogen flame, emits brilliant white light (known as limelight).
- When exposed to atmosphere,  
 $\text{CaO} + \text{H}_2\text{O} \longrightarrow \text{Ca(OH)}_2$   
(Moisture) (Slaked Lime)
- $\text{CaO} + \text{CO}_2 \longrightarrow \text{CaCO}_3$   
(Limestone)
- Being a basic oxide, can combine with acidic impurities.  
 $\text{CaO} + \text{SiO}_2 \xrightarrow{\Delta} \text{CaSiO}_3$

#### Uses :

- Purification of sugar.
- Important constituent of cement, mortar, caustic soda lime ( $\text{CaO} + \text{NaOH}$ ) etc.

### Calcium Carbonate (Limestone) ; $\text{CaCO}_3$

#### Preparation :

- From slaked lime :  
 $\text{Ca(OH)}_2 + \text{CO}_2 \longrightarrow \text{CaCO}_3 + \text{H}_2\text{O}$
- From calcium chloride :  
 $\text{CaCl}_2 + \text{Na}_2\text{CO}_3 \longrightarrow \text{CaCO}_3 + 2\text{NaCl}$

#### Properties :

- White fluffy solid and insoluble in water.
- $\text{CaCO}_3 \xrightarrow{\Delta} \text{CaO} + \text{CO}_2$
- Liberates  $\text{CO}_2$   
 $\text{CaCO}_3 + 2\text{HCl} \longrightarrow \text{CaCl}_2 + \text{CO}_2\uparrow + \text{H}_2\text{O}$   
 $\text{CaCO}_3 + \text{H}_2\text{SO}_4 \longrightarrow \text{CaSO}_4 + \text{CO}_2\uparrow + \text{H}_2\text{O}$

#### Uses :

- In building material, in the manufacturing of quick lime,  $\text{Na}_2\text{CO}_3$ .
- In the extraction of metals.
- As an antacid, abrasive in toothpaste, filler in cosmetics.

### Calcium Sulphate (Plaster of Paris) ; $(\text{CaSO}_4)_2 \cdot \text{H}_2\text{O}$

Plaster of Paris is calcium sulphate hemihydrate  
 $\text{CaSO}_4 \cdot \frac{1}{2} \text{H}_2\text{O}$ .

#### Preparation :

- $2(\text{CaSO}_4 \cdot 2\text{H}_2\text{O}) \xrightarrow[393 \text{ K}]{\Delta} 2(\text{CaSO}_4) \cdot \text{H}_2\text{O} + 3\text{H}_2\text{O}$   
Gypsum Plaster of Paris
- $2(\text{CaSO}_4) \cdot \text{H}_2\text{O} \xrightarrow[473 \text{ K}]{\Delta} 2\text{CaSO}_4 + \text{H}_2\text{O}$   
(Dead burnt plaster)

#### Properties :

- It is white powder.
- On mixing with water, it sets into hard mass after sometime.  
 $(\text{CaSO}_4)_2 \cdot \text{H}_2\text{O} + 3\text{H}_2\text{O} \longrightarrow 2(\text{CaSO}_4 \cdot 2\text{H}_2\text{O})$   
Gypsum

#### Uses :

- For producing moulds for pottery, ceramic etc.
- For making statues, models and other decorative materials.
- In surgical bandages.
- In dentistry.

### Cement

It was first introduced in England in 1824 by Joseph Aspdin. It is also called Portland cement because it resembles with the famous building stone found near Portland in England. It is a finely ground mixture of calcium silicates and aluminates which set to a hard mass when treated with water.

#### Composition of Cement

$\text{CaO} = 50\text{-}60\%$	$\text{MgO} = 2\text{-}3\%$
$\text{SiO}_2 = 20\text{-}25\%$	$\text{Fe}_2\text{O}_3 = 1\text{-}2\%$
$\text{Al}_2\text{O}_3 = 5\text{-}10\%$	$\text{SO}_3 = 1\text{-}2\%$

# INFOSHOTS

## Sodium is associated with an increased risk of type 2 diabetes!

According to a latest research, sodium intake may be linked to an increased risk of developing both type 2 diabetes (T2D) and Latent Autoimmune Diabetes in Adults (LADA).

The main source of sodium in the diet is through salt. Salt (sodium chloride) contains 40% sodium, thus for every 2.5 g of salt consumed, 1 g is sodium. Research has suggested that excessive salt consumption may increase the risk of developing T2D, possibly through a direct effect on insulin resistance, and/or by promoting high blood pressure and weight gain.

## Biological Importance

### Sodium and Potassium

$\text{Na}^+$  :

- Major component of blood plasma.
- Helps in nerve signal transmission and function of heart.
- It maintains the electrolyte balance in the body.
- In transport of sugar and amino acids into cell.

$\text{K}^+$  :

- Activating some enzymes.
- Produces energy rich ATP molecules.
- Protein synthesis.
- For transmission of nerve signals.

### Calcium and Magnesium

$\text{Ca}^{2+}$  :

- In stabilisation of protein.
- As a messenger for hormonal action.
- As a trigger for the contraction of muscles.
- In maintenance of rhythm of heart.

$\text{Mg}^{2+}$  :

- Catalyses enzymatic reactions.
- In photosynthesis.
- With ATP forms complex, which release energy.

# SPEED PRACTICE

- In the absence of hydrogen bonding, the boiling point of water would have been  
(a)  $100^\circ\text{C}$  (b)  $0^\circ\text{C}$  (c)  $-80^\circ\text{C}$  (d)  $373^\circ\text{C}$
- When sodium reacts with sufficient amount of air, the product obtained is  
(a)  $\text{Na}_2\text{O}$  (b)  $\text{Na}_2\text{O}_2$  (c)  $\text{NaO}_2$  (d)  $\text{NaO}$
- Identify the incorrect statement about heavy water.  
(a) It reacts with  $\text{SO}_3$  to form deuterated sulphuric acid ( $\text{D}_2\text{SO}_4$ ).  
(b) It is used as a coolant in nuclear reactors.  
(c) It reacts with  $\text{CaC}_2$  to produce  $\text{C}_2\text{D}_2$  and  $\text{Ca(OD)}_2$ .  
(d) It reacts with  $\text{Al}_4\text{C}_3$  to produce  $\text{CD}_4$  and  $\text{Al(OD)}_3$ .
- Which of the following complex is the most stable?  
(a)  $[\text{Be}(\text{H}_2\text{O})_4]^{2+}$  (b)  $[\text{Mg}(\text{H}_2\text{O})_4]^{2+}$   
(c)  $[\text{Ca}(\text{H}_2\text{O})_4]^{2+}$  (d)  $[\text{Sr}(\text{H}_2\text{O})_4]^{2+}$


(JEE Main Online 2016)






5.  $K_2CO_3$  cannot be prepared by Solvay process because  
 (a)  $K_2CO_3$  decomposes in the process  
 (b)  $KHCO_3$  is soluble in water  
 (c)  $KHCO_3$  is insoluble in water  
 (d)  $KHCO_3$  is thermally stable.
6. A metal  $M$  readily forms a water-soluble sulphate  $MSO_4$ , a water-insoluble hydroxide  $M(OH)_2$  and an oxide  $MO$  which becomes inert on heating. The hydroxide is soluble in  $NaOH$ . The metal  $M$  is  
 (a) Be (b) Mg (c) Ca (d) Sr
7. Ionic metal hydrides react with water to give hydrogen gas. Which of the following hydrides will produce maximum amount of  $H_2$  if equal mass of each hydride is taken?  
 (a)  $LiH$  (b)  $CaH_2$  (c)  $NaH$  (d)  $MgH_2$
8. Both lithium and magnesium display several similar properties due to the diagonal relationship, however, the one which is incorrect, is  
 (a) both form nitrides  
 (b) nitrates of both  $Li$  and  $Mg$  yield  $NO_2$  and  $O_2$  on heating  
 (c) both form basic carbonates  
 (d) both form soluble bicarbonates.
- (JEE Main 2017)
9. The hydride ion,  $H^-$  is a stronger base than hydroxide ion,  $OH^-$ . Which of the following reactions will occur if sodium hydride ( $NaH$ ) is dissolved in water?  
 (a)  $H^-_{(aq)} + H_2O_{(l)} \longrightarrow H_3O^+_{(aq)}$   
 (b)  $H^-_{(aq)} + H_2O_{(l)} \longrightarrow OH^-_{(aq)} + H_{2(g)}$   
 (c)  $H^-_{(aq)} + H_2O_{(l)} \longrightarrow \text{No reaction}$   
 (d)  $H^-_{(aq)} + 2H_2O_{(l)} \longrightarrow H_3O^+_{(aq)} + H_{2(g)}$
10. What is the product obtained when  $MgCl_2 \cdot 6H_2O$  is strongly heated?  
 (a)  $MgCl_2$  (b)  $MgO$   
 (c)  $MgCl_2 \cdot 2H_2O$  (d)  $MgCl_2 \cdot 4H_2O$
11. Which of the following hydrides is not ionic?  
 (a)  $CaH_2$  (b)  $BaH_2$  (c)  $SrH_2$  (d)  $BeH_2$
12. The compound formed when gypsum is dissolved in aqueous ammonium sulphate solution is  
 (a)  $CaSO_4 \cdot NH_4Cl \cdot H_2O$   
 (b)  $CaCl_2(NH_4)_2SO_4 \cdot H_2O$   
 (c)  $CaSO_4(NH_4)_2SO_4 \cdot 2H_2O$   
 (d)  $CaCl_2 \cdot NH_4Cl \cdot 2H_2O$
13. Which of the following statements about hydrogen is incorrect?  
 (a) Hydronium ion,  $H_3O^+$  exists freely in solution.  
 (b) Dihydrogen does not act as a reducing agent.  
 (c) Hydrogen has three isotopes of which tritium is the most common.  
 (d) Hydrogen never acts as cation in ionic salts.
- (NEET Phase-I 2016)
14. The solubility of hydroxides, fluorides or oxalates of the metals of group II  
 (a) increases down the group  
 (b) decreases down the group  
 (c) varies randomly  
 (d) is same.
15. Polyphosphates are used as water softening agents because they  
 (a) form soluble complexes with anionic species  
 (b) form soluble complexes with cationic species  
 (c) precipitate cationic species  
 (d) precipitate anionic species.
16. The correct sequence of increasing covalent character is represented by  
 (a)  $LiCl < NaCl < BeCl_2$  (b)  $BeCl_2 < LiCl < NaCl$   
 (c)  $NaCl < LiCl < BeCl_2$  (d)  $BeCl_2 < NaCl < LiCl$
17. A 'hair-bleach' solution contains  $H_2O_2$ . The amount of  $H_2O_2$  in a 12.5 g sample was determined by titration with  $KMnO_4$  in acidic medium. 39.3 mL of 0.5045 N  $KMnO_4$  was required by the sample. What is the mass percentage of  $H_2O_2$  in the bleach solution?  
 (a) 2.69 (b) 3.15 (c) 3.37 (d) 2.12
18. Ionic mobility of which of the following alkali metal ions is lowest when aqueous solution of their salts are put under an electric field?  
 (a) K (b) Rb  
 (c) Li (d) Na
- (NEET 2017)
19. When a substance ( $A$ ) reacts with water it produces a combustible gas ( $B$ ) and a solution of substance ( $C$ ) in water. When another substance ( $D$ ) reacts with this solution of ( $C$ ) it produces the same gas ( $B$ ) on warming. ( $D$ ) may also produce the gas ( $B$ ) on reaction with dilute sulphuric acid at room temperature. ( $A$ ) imparts a deep golden yellow colour to a smokeless flame of Bunsen burner.  $A$ ,  $B$ ,  $C$  and  $D$  are respectively







13. (b, c) : Dihydrogen acts as a powerful reducing agent and reduces metal oxides such as CuO, ZnO, PbO and Fe<sub>3</sub>O<sub>4</sub> to their respective metals.



Hydrogen has three isotopes of which protium is the most common and tritium is radioactive.

14. (a) : Both lattice energy and hydration energy decrease down the group due to gradual increase in size of  $M^{2+}$  ion. Lattice energy tends to increase the solubility while hydration energy tends to decrease the solubility. However, lattice energy predominates hydration energy. Therefore, solubility increases down the group.

15. (b) : Polyphosphates, e.g., calgon are used as water softening agents because they form soluble complexes with cations,  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ .

16. (c) : LiCl and BeCl<sub>2</sub> are predominantly covalent compounds but NaCl is an ionic compound. The covalent character of LiCl and BeCl<sub>2</sub> is due to polarisation of Cl<sup>-</sup> ion by Li<sup>+</sup> and Be<sup>2+</sup> ions. As the polarisation is more due to Be<sup>2+</sup> ions as it has smaller size and greater charge, BeCl<sub>2</sub> is more covalent than LiCl.


17. (a) :  $2\text{MnO}_4^- + 5\text{H}_2\text{O}_2 + 6\text{H}^+ \longrightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{O}_2$   
1000 mL of 1N  $\text{MnO}_4^- \equiv 1000 \text{ mL of } 1\text{N } \text{H}_2\text{O}_2$

$$\begin{aligned} \text{Equivalent of } \text{H}_2\text{O}_2 &= \text{equivalent of } \text{MnO}_4^- \\ &= \frac{39.3 \times 0.5045}{1000} = 0.0198 \end{aligned}$$


$$\therefore \text{Amount of pure } \text{H}_2\text{O}_2 = 0.0198 \times 17 = 0.337 \text{ g}$$

$$\% \text{ of } \text{H}_2\text{O}_2 \text{ in bleach solution} = \frac{0.337}{12.5} \times 100 = 2.69 \%$$



18. (c) : The hydration enthalpy of alkali metal ions decreases with increase in ionic sizes i.e.,




Hence, lithium having maximum degree of hydration, will be least mobile. Thus, the order of ionic mobility is :



19. (a) :  $2\text{Na} + 2\text{H}_2\text{O} \longrightarrow 2\text{NaOH} + \text{H}_2$   
(A) (C) (B)



20. (c) : Thermal stability of carbonates of alkaline earth metals increases with increase in atomic mass down the group or with increasing cationic size :



21. (b) : Amount of  $\text{MgSO}_4 = 0.002 \times 120 = 0.24 \text{ g of } \text{MgSO}_4$

$$\text{Also, } 120 \text{ g of } \text{MgSO}_4 \equiv 100 \text{ g of } \text{CaCO}_3$$

$$\therefore 0.24 \text{ g of } \text{MgSO}_4 \equiv \frac{100 \times 0.24}{120} = 0.2 \text{ g of } \text{CaCO}_3$$

$$1 \text{ L of water} = 10^3 \text{ mL} = 1000 \text{ g } (\because \rho \text{ H}_2\text{O} = 1 \text{ g/cc})$$


$$\text{Thus, } 10^3 \text{ g of water contains} = 0.2 \text{ g of } \text{CaCO}_3$$

Hence,  $10^6 \text{ g of water}$  shall contain

$$= \frac{0.2 \times 10^6}{10^3} = 200 \text{ g}$$

Therefore, hardness of water is 200 ppm.

22. (a) : All elements of s-block (group I and group II) except Be and Mg give characteristic colour to the flame.



22400 mL of O<sub>2</sub> is formed from 68 g of H<sub>2</sub>O<sub>2</sub>.

$$\therefore 10 \text{ mL of O}_2 \text{ is formed by } \frac{68}{22400} \times 10 \text{ g of } \text{H}_2\text{O}_2 = 0.0303 \text{ g H}_2\text{O}_2$$

$$1 \text{ mL of } \text{H}_2\text{O}_2 \text{ solution contains } 0.0303 \text{ g H}_2\text{O}_2$$

$$\therefore 100 \text{ mL of } \text{H}_2\text{O}_2 \text{ solution contains} = 0.0303 \times 100 = 3.035 \text{ g} = 3.035\% \text{ of } \text{H}_2\text{O}_2$$

24. (c) : Beryllium salts are readily hydrolysed.

25. (b) : Water molecule can form two hydrogen bonds through its two hydrogen atoms and another two hydrogen bond with two lone pairs of electrons of O-atom.

26. (c)

27. (c) : Permanent hardness of water is due to the presence of chlorides and sulphates of calcium and magnesium.


28. (b) : KOH is better absorber of CO<sub>2</sub> because K<sub>2</sub>CO<sub>3</sub> formed is more soluble and does not separate out.

29. (a) : Lithium does not form peroxide.

30. (c) :  $\text{CaOCl}_2 + \text{CO}_2 \longrightarrow \text{CaCO}_3 + \text{Cl}_2$



# EXAMINER'S MIND CLASS XI



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XI.  
This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.


<b>Section - I</b>	Q. 1 to 10 Only One Option Correct Type MCQs.
<b>Section - II</b>	Q. 11 to 13 More than One Options Correct Type MCQs.
<b>Section - III</b>	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
<b>Section - IV</b>	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
<b>Section - V</b>	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
<b>Section - VI</b>	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).

## HYDROCARBONS

### SECTION - I



#### Only One Option Correct Type

1. The electrophile,  $E^+$  attacks the benzene ring to generate the intermediate  $\sigma$ -complex. Which of the following,  $\sigma$ -complex is of the lowest energy?




2. 3-Octyne is synthesized by adding a bromoalkane in a mixture of sodium amide and an alkyne. The bromoalkane and alkyne are respectively
- (a)  $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$  and  $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CH}$   
 (b)  $\text{BrCH}_2\text{CH}_2\text{CH}_3$  and  $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$   
 (c)  $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$  and  $\text{CH}_3\text{C}\equiv\text{CH}$   
 (d)  $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$  and  $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CH}$

3. *trans*-2-Phenyl-1-bromocyclopentane on reaction with alcoholic KOH produces
- (a) 4-phenylcyclopentene  
 (b) 2-phenylcyclopentene  
 (c) 1-phenylcyclopentene  
 (d) 3-phenylcyclopentene.
4. When cyclohexane is poured on water, it floats, because
- (a) cyclohexane is in 'boat' form  
 (b) cyclohexane in 'chair' form is heavier than water  
 (c) cyclohexane is more dense than water  
 (d) cyclohexane is less dense than water.
5. Which of the compound with molecular formula  $\text{C}_5\text{H}_{10}$  yields acetone on ozonolysis?
- (a) 3-Methyl-1-butene  
 (b) Cyclopentane  
 (c) 2-Methyl-1-butene  
 (d) 2-Methyl-2-butene
6. The *Z*-isomer among the following is
- (a)







7. Which of the given reaction conditions will give maximum yields of  $\text{C}_2\text{H}_5\text{Cl}$ ?


- (a)  $\text{C}_2\text{H}_6$  (excess) +  $\text{Cl}_2 \xrightarrow{\text{UV light}}$   
 (b)  $\text{C}_2\text{H}_6 + \text{Cl}_2 \xrightarrow[\text{room temperature}]{\text{dark}}$   
 (c)  $\text{C}_2\text{H}_6 + \text{Cl}_2$  (excess)  $\xrightarrow{\text{UV light}}$   
 (d)  $\text{C}_2\text{H}_6 + \text{Cl}_2 \xrightarrow{\text{UV light}}$





The major products P and Q are

- (a)  and  $\text{CH}_3\text{CH}_2\text{CHO}$   
 (b)  and  $\text{CH}_3\text{COCH}_3$   
 (c)  and  $\text{CH}_3\text{COCH}_3$   
 (d)  and  $\text{CH}_3\text{CH}_2\text{CHO}$

9. In the following reaction,



- (a)  $\text{Me}-\text{CH}=\text{CH}_2$  (b)  $\text{CH}_2=\text{CH}_2$   
 (c)  (d) 



Identify product P.


- (a)  $\text{CH}_3-\text{CH}(\text{Cl})-\text{CH}_2\text{NO}$  (b)  $\text{CH}_3-\text{CH}(\text{NO})-\text{CH}_2\text{Cl}$   
 (c)  $\text{CH}_3-\text{CH}_2-\text{CH}(\text{Cl})\text{NO}$  (d)  $\text{CH}_2=\text{CH}-\text{CH}_2\text{Cl}$

## SECTION - II

More than One Options Correct Type

11. Which of the following are cumulative dienes?


- (a)  $\text{CH}_3-\text{CH}=\text{C}=\text{CH}-\text{CH}_3$   
 (b)  $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$   
 (c)  $\text{CH}_2=\text{CH}-\text{CH}_2\text{CH}=\text{CH}_2$   
 (d)  $\text{CH}_3\text{CH}_2\text{CH}=\text{C}=\text{CH}-\text{CH}_3$



Products of the given reaction are

- (a)  $\text{CH}_3-\text{C}(\text{CH}_3)(\text{OH})-\text{CH}_3$  (b)  $\text{HOCH}_2-\text{C}(\text{CH}_3)_2-\text{CH}_3$   
 (c)  $\text{CH}_3-\text{C}(\text{CH}_3)(\text{OH})-\text{CH}_2-\text{CH}=\text{CH}_2$   
 (d)  $\text{CH}_3-\text{C}(\text{CH}_3)(\text{OH})-\text{CH}=\text{CH}-\text{CH}_3$

13. Among P, Q, R and S the aromatic compounds are




## SECTION - III

Paragraph Type

### Paragraph for Questions 14 and 15

An acyclic hydrocarbon P, having molecular formula  $\text{C}_6\text{H}_{10}$  gave acetone as the only organic product through the following sequence of reactions in which Q is an intermediate organic compound.






14. The structure of the compound P is

- (a)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$   
 (b)  $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_3$   
 (c)  $\begin{array}{c} \text{H}_3\text{C} \\ \diagup \\ \text{H}-\text{C}-\text{C}\equiv\text{C}-\text{CH}_3 \\ \diagdown \\ \text{H}_3\text{C} \end{array}$   
 (d)  $\begin{array}{c} \text{H}_3\text{C} \\ \diagup \\ \text{H}_3\text{C}-\text{C}-\text{C}\equiv\text{C}-\text{H} \\ \diagdown \\ \text{H}_3\text{C} \end{array}$

15. The structure of the compound Q is

- (a)  $\begin{array}{c} \text{H}_3\text{C} \\ \diagup \\ \text{H}-\text{C}-\text{C}-\text{CH}_2\text{CH}_3 \\ \diagdown \\ \text{H}_3\text{C} \\ | \\ \text{H} \end{array}$   
 (b)  $\begin{array}{c} \text{H}_3\text{C} \\ \diagup \\ \text{H}_3\text{C}-\text{C}-\text{C}-\text{CH}_3 \\ \diagdown \\ \text{H}_3\text{C} \\ | \\ \text{H} \end{array}$   
 (c)  $\begin{array}{c} \text{H}_3\text{C} \\ \diagup \\ \text{H}-\text{C}-\text{CH}_2-\text{CHCH}_3 \\ \diagdown \\ \text{H}_3\text{C} \\ | \\ \text{OH} \end{array}$   
 (d)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$

#### Paragraph for Questions 16 and 17



To evaluate resonance energy of benzene, we can compare heat of hydrogenation of cyclohexene and benzene (or its derivative).

16. Heat of hydrogenation of cyclohexene is

- (a)  $-85.8 \text{ kcal mol}^{-1}$   
 (b)  $-49.8 \text{ kcal mol}^{-1}$   
 (c)  $-36.0 \text{ kcal mol}^{-1}$   
 (d)  $-28.6 \text{ kcal mol}^{-1}$

17. Resonance energy of benzene is

- (a)  $-49.8 \text{ kcal mol}^{-1}$   
 (b)  $-36 \text{ kcal mol}^{-1}$   
 (c)  $+36.0 \text{ kcal mol}^{-1}$   
 (d)  $-28.6 \text{ kcal mol}^{-1}$

#### SECTION - IV

##### Matching List Type

18. Match the list I with list II and select the correct answer using the codes given below the lists :

##### List I

- (P) Benzene  
 (Q) Ethylene  
 (R) Acetaldehyde  
 (S) Chloroform

##### List II

- (i) Phosgene  
 (ii) Silver mirror test  
 (iii) Mustard gas  
 (iv)  $(4n + 2)\pi$  electrons  
 (v) Carbylamine

- |     | P    | Q     | R    | S     |
|-----|------|-------|------|-------|
| (a) | (iv) | (iii) | (ii) | (i)   |
| (b) | (v)  | (ii)  | (i)  | (iv)  |
| (c) | (ii) | (iv)  | (i)  | (iii) |
| (d) | (v)  | (i)   | (iv) | (iii) |

19. Match the list I with list II and select the correct answer using the codes given below the lists :

##### List I

- (P) Alkyl halide + Sodium in presence of dry ether  
 (Q) Arene + Acid halide in presence of anhyd.  $\text{AlCl}_3$   
 (R) Arene + Fuming sulphuric acid  
 (S) Arene + Hydrogen in presence of Ni

##### List II

- (i) Sulphonation  
 (ii) Wurtz reaction  
 (iii) Catalytic hydrogenation  
 (iv) Friedel-Crafts reaction

- |     | P     | Q     | R     | S     |
|-----|-------|-------|-------|-------|
| (a) | (i)   | (iii) | (ii)  | (iv)  |
| (b) | (iv)  | (ii)  | (iii) | (i)   |
| (c) | (iii) | (i)   | (iv)  | (ii)  |
| (d) | (ii)  | (iv)  | (i)   | (iii) |

#### SECTION - V

##### Assertion Reason Type

20. **Assertion :** 1-butene on reaction with HBr in presence of peroxide produces 1-bromobutane.

**Reason :** It involves the formation of a primary radical.

21. **Assertion :** Benzene on heating with conc.  $\text{H}_2\text{SO}_4$  gives benzenesulphonic acid which when heated with superheated steam under pressure gives benzene.

**Reason :** Sulphonation is a reversible process.


22. **Assertion :** The gauche conformation of ethylene glycol is more stable than its anti-conformation.

**Reason :** The gauche conformation is stabilized by H-bonding.

## SECTION - VI

### Integer Value Correct Type

23. The number of optically active products obtained from the complete ozonolysis of the given compound is



24. The total number of alkenes possible by dehydrobromination of 3-bromo-3-cyclopentylhexane using alcoholic KOH is

25. Which of the following molecules have zero dipole moment?


*cis*-1, 2-Dichloroethene, *trans*-1, 2-dichloroethene, 1, 1-dichloroethene, *trans*-2-pentene, *cis*-2-pentene, 1-butyne, 2-butyne, *trans*-2-butene, *cis*-2-butene

## THERMODYNAMICS

### SECTION - I


#### Only One Option Correct Type

1. For which of the following process,  $q = \Delta U$ ?



- (a)  $O \rightarrow A$  (b)  $O \rightarrow D$   
(c)  $O \rightarrow B$  (d)  $O \rightarrow C$
2. Standard entropies of  $X_2$ ,  $Y_2$  and  $XY_3$  are 60, 40 and  $50 \text{ J K}^{-1} \text{ mol}^{-1}$  respectively. For the reaction,  $\frac{1}{2} X_2 + \frac{3}{2} Y_2 \rightarrow XY_3$ ,  $\Delta H^\circ = -30 \text{ kJ mol}^{-1}$ , to be at equilibrium, the temperature will be  
(a) 500 K (b) 750 K (c) 1000 K (d) 1250 K
3. Internal energy and pressure for one mole of a gas of unit volume are related as  
(a)  $p = \frac{2}{3} E$  (b)  $p = \frac{E}{2}$  (c)  $p = \frac{3}{2} E$  (d)  $p = 2E$
4. One mole of a non-ideal gas undergoes a change of state (2.0 atm, 3.0 L, 95 K)  $\rightarrow$  (4.0 atm, 5.0 L, 245 K) with a change in internal energy,  $\Delta U = 30.0 \text{ L atm}$ . The change in enthalpy ( $\Delta H$ ) of the process in L atm is  
(a) 40.0 (b) 42.3  
(c) 44.0  
(d) not defined, because pressure is not constant.

5. The direct conversion of A to B is difficult, hence it is carried out by the following path :






Entropy change ( $\Delta S$ ) for the process,

$A \rightarrow C = 50 \text{ e.u.}$  ;  $C \rightarrow D = 30 \text{ e.u.}$  ;  $B \rightarrow D = 20 \text{ e.u.}$  (where e.u. is unit of entropy), then  $\Delta S(A \rightarrow B)$  is

- (a) +100 e.u. (b) +60 e.u.  
(c) -100 e.u. (d) -60 e.u.

6. Which of the algebraic relationships on the basis of given thermochemical equations is correct?




- (a)  $z = x + y$  (b)  $x = y - z$   
(c)  $x = y + z$  (d)  $y = 2z - x$

7. An athlete has taken 100 g glucose of energy equivalent to 1560 kJ. He utilizes 50% of this gained energy in an event. In order to avoid storage of energy in the body, calculate the mass of water he would need to perspire. (Enthalpy of  $\text{H}_2\text{O}$  for evaporation =  $44 \text{ kJ mol}^{-1}$ )

- (a) 346 g (b) 316 g (c) 323 g (d) 319 g

8. The volume of a gas is reduced to half from its original volume. Its specific heat will be  
(a) reduced to half (b) doubled  
(c) remain constant (d) increased four times.

9. Acetic acid,  $\text{CH}_3\text{COOH}$  can form a dimer  $(\text{CH}_3\text{COOH})_2$  in the gas phase. The dimer is held together by two H-bonds with a total strength of 66.5 kJ per mol of dimer.



If at  $25^\circ \text{C}$ , the equilibrium constant for the dimerization is  $1.3 \times 10^3$  then  $\Delta S^\circ$  for the reaction is

- (a) 0.163 kJ (b) 0.2010 kJ  
(c) 0.1251 kJ (d) 0.091 kJ.

10. Standard enthalpies of formation of  $O_3$ ,  $CO_2$ ,  $NH_3$  and  $HI$  are  $+142.2$ ,  $-393.2$ ,  $-46.2$  and  $+25.9 \text{ kJ mol}^{-1}$  respectively. The order of their increasing stabilities will be
- (a)  $O_3$ ,  $CO_2$ ,  $NH_3$ ,  $HI$  (b)  $CO_2$ ,  $NH_3$ ,  $HI$ ,  $O_3$   
 (c)  $O_3$ ,  $HI$ ,  $NH_3$ ,  $CO_2$  (d)  $NH_3$ ,  $HI$ ,  $CO_2$ ,  $O_3$

### SECTION - II

#### More than One Options Correct Type


11. Select the correct option for the equilibrium under standard conditions as given :
- $$H_2O_{(s)} \rightleftharpoons H_2O_{(l)} ; \Delta S_1^\circ$$
- $$H_2O_{(l)} \rightleftharpoons H_2O_{(v)} ; \Delta S_2^\circ$$
- $$H_2O_{(s)} \rightleftharpoons H_2O_{(v)} ; \Delta S_3^\circ$$
- (a)  $\Delta S_1^\circ > \Delta S_2^\circ$  (b)  $\Delta S_2^\circ \gg \Delta S_1^\circ$   
 (c)  $\Delta S_3^\circ > \Delta S_2^\circ$  (d)  $\Delta S_3^\circ > \Delta S_1^\circ$
12. Which of the following statements are not correct?
- (a) The presence of reacting species in a covered beaker is an example of open system.  
 (b) There is an exchange of energy as well as matter between the system and the surroundings in a closed system.  
 (c) The presence of reactants in a closed vessel made up of copper is an example of a closed system.  
 (d) The presence of reactants in a thermos flask or any other closed insulated vessel is an example of a closed system.
13. Which of the following expressions regarding entropy are correct?
- (a)  $\Delta S_{\text{system}} = \frac{q}{T}$   
 (b)  $\Delta S_{\text{system}} = \Delta S_{\text{total}} + \Delta S_{\text{surrounding}}$   
 (c)  $\Delta S = S_{\text{final}} - S_{\text{initial}}$   
 (d)  $\Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surrounding}}$

### SECTION - III

#### Paragraph Type

#### Paragraph for Questions 14 and 15

A sample containing 1 mole of a monoatomic ideal gas ( $C_v = \frac{3}{2}R$ ) is taken through the cycle as shown :




14. Temperature at points 1, 2 and 3 are, respectively  
 (a) 273 K, 546 K, 273 K (b) 546 K, 273 K, 273 K  
 (c) 273 K, 273 K, 273 K (d) 546 K, 546 K, 273 K.
15.  $\Delta E$  for the process (1  $\rightarrow$  2) is  
 (a) 3.4 J (b)  $3.4 \times 10^3 \text{ J}$   
 (c)  $-3.4 \text{ J}$  (d)  $-3.4 \times 10^3 \text{ J}$ .

#### Paragraph for Questions 16 and 17

Entropy is a measure of degree of randomness. Entropy is directly proportional to temperature. Every system tries to acquire state of maximum randomness or disorder. Entropy is a measure of unavailable energy. Unavailable energy = Entropy  $\times$  Temperature. The ratio of entropy of vapourisation and boiling point of a substance remains almost constant.

16. Observe the graph and identify the correct statement.



- (a)  $T_1$  is melting point and  $T_2$  is boiling point.  
 (b)  $T_1$  is boiling point and  $T_2$  is melting point.  
 (c)  $\Delta S_{\text{fusion}}$  is more than  $\Delta S_{\text{vap}}$ .  
 (d)  $T_2$  is lower than  $T_1$ .
17. The law of thermodynamics formulated by Nernst, which helps to determine absolute entropy is  
 (a) zeroth law (b) first law  
 (c) second law (d) third law.

### SECTION - IV

#### Matching List Type

18. Match the list I with list II and select the correct answer using the codes given below the list :

List I	List II
(P) $H_2 \rightarrow 2H$	1. $\Delta H = B.E. (C \equiv C) + 2B.E. (H-H) - B.E. (C-C) - 4B.E. (C-H)$
(Q) $CH_4 \rightarrow C + 4H$	2. $\Delta H = \Delta_f H \text{ of } H \times 2$
(R) $C_2H_2 + 2H_2 \rightarrow C_2H_6$	3. $\Delta H = \text{Average bond energy} \times 4$
(S) $C_2H_4 + H_2 \rightarrow C_2H_6$	4. $\Delta H = B.E.(C=C) + B.E. (H-H) - B.E. (C-C) - 2B.E. (C-H)$



	P	Q	R	S
(a)	1	2	3	4
(b)	2	3	1	4
(c)	3	2	4	1
(d)	1	3	4	2

19. Match the list I with list II and select the correct answer using the codes given below the list :


List I	List II
(P) $\text{H}_{2(g)} + \text{Br}_{2(g)} \rightarrow 2\text{HBr}_{(g)}$	1. $\Delta H = \Delta U - 2RT$
(Q) $\text{PCl}_{5(g)} \rightarrow \text{PCl}_{3(g)} + \text{Cl}_{2(g)}$	2. $\Delta H = \Delta U + 3RT$
(R) $\text{N}_{2(g)} + 3\text{H}_{2(g)} \rightarrow 2\text{NH}_{3(g)}$	3. $\Delta H = \Delta U$
(S) $2\text{N}_2\text{O}_{5(g)} \rightarrow 4\text{NO}_{2(g)} + \text{O}_{2(g)}$	4. $\Delta H = \Delta U + RT$

P	Q	R	S
(a)	3	1	2
(b)	3	4	1
(c)	2	1	4
(d)	4	2	1

### SECTION - V

#### Assertion Reason Type

20. **Assertion :** The isothermal reversible work of expansion is always greater than that of irreversible expansion of an ideal gas.  
**Reason :** Work is a path dependent function.
21. **Assertion :** The entropy change in the reaction.



**Reason :** Entropy increases when number of gaseous particles (*i.e.*, molecules or ions) of the product increases.

22. **Assertion :** If both  $\Delta H^\circ$  and  $\Delta S^\circ$  are positive then reaction will be spontaneous at high temperature.  
**Reason :** All processes with positive entropy change are spontaneous.

### SECTION - VI

#### Integer Value Correct Type


23. The freezing point of *iso*-butane is  $-160^\circ\text{C}$ .  $\Delta H_{(\text{solid} \rightarrow \text{liquid})}$  is  $+4520 \text{ J mol}^{-1}$ . For this fusion process, entropy change in  $\text{J mol}^{-1}$  is  $10y$ . The value of  $y$  is
24. At dynamic equilibrium, the reaction on both sides occur at the same rate and the mass on both sides of the equilibrium does not undergo any change. This condition can be achieved only when the value of  $\Delta G$  is
25. In a constant volume calorimeter, 3.5 g of a gas with molecular weight 28, was burnt in excess of oxygen at 298 K. The temperature of the calorimeter was found to increase from 298 K to 298.45 K due to the combustion process. Given that, heat capacity of the calorimeter is  $2.5 \text{ kJ K}^{-1}$ , the numerical value for the enthalpy of combustion of the gas in  $\text{kJ mol}^{-1}$  is

## SOLUTIONS


### HYDROCARBONS

1. (b):  $-\text{NO}_2$  is an electron-withdrawing group and hence, will destabilise  $\sigma$ -complex thus, others are of higher energy.

2. (d):




3. (d):




4. (d)                      5. (d)                      6. (a)


7. (a):  $\text{C}_2\text{H}_6$  must be in excess for the best yield of  $\text{C}_2\text{H}_5\text{Cl}$ .

8. (c): It is cumene hydroperoxide rearrangement reaction.






In Hofmann elimination reaction, less sterically hindered  $\beta$ -hydrogen is removed and less substituted alkene is the major product.




11. (a, d)

12. (c, d)

13. (a, b, c, d):



14. (d): The formula,  $\text{C}_6\text{H}_{10}$  resemble  $\text{C}_n\text{H}_{2n-2}$  which shows P is an alkyne.



15. (b)

16. (d): Cyclohexene has one  $\text{C}=\text{C}$  bond.





$$\Delta H \text{ of cyclohexene} = -85.8/3 = -28.6 \text{ kcal mol}^{-1}$$

17. (c)

18. (a)


19. (d)

20. (c): Formation of 1-bromobutane from 1-butene involves secondary free radical as intermediate.




21. (a)

22. (a)



24. (5):



25. (3): *trans*-1,2-Dichloroethene, *trans*-2-butene, 2-butyne have zero dipole moment.

# BEST TOOLS FOR SUCCESS IN JEE Main



**10** Very Similar Practice Test Papers

**16** Years JEE MAIN 2017-2015(Offline & Online)-2013 & AIEEE (2012-2002)



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email: info@mtg.in

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!




## THERMODYNAMICS

1. (b):  $w = -P\Delta V$

During process  $O \rightarrow D$ , no change in volume *i.e.*,  $\Delta V = 0$

$$\therefore w = -P\Delta V = 0; \Delta U = q - w \Rightarrow \Delta U = q$$

2. (b): For the reaction,



$$\Delta S^\circ = \Sigma S^\circ_{(\text{products})} - \Sigma S^\circ_{(\text{reactants})}$$

$$\Delta S^\circ = 50 - \left[ 40 \times \frac{3}{2} + 60 \times \frac{1}{2} \right] = -40 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\therefore \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

At equilibrium,  $\Delta G^\circ = 0$

$$\therefore \Delta H^\circ = T\Delta S^\circ$$

$$T = \frac{\Delta H^\circ}{\Delta S^\circ} = \frac{-30 \times 10^3}{-40} = 750 \text{ K}$$

3. (a):  $\therefore PV = RT$  (For one mole)

At unit volume;  $p \times 1 = RT$

$$\text{Internal energy, } E = \frac{3}{2}RT \Rightarrow \frac{3}{2}p$$

$$\therefore p = \frac{2}{3}E$$

4. (c):  $\Delta H = H_2 - H_1 = (E_2 + P_2V_2) - (E_1 + P_1V_1)$   
 $= (E_2 - E_1) + (P_2V_2 - P_1V_1)$   
 $= 30 + (4 \times 5 - 2 \times 3) = 44 \text{ L atm}$

5. (b):  $\Delta S(A \rightarrow B) = \Delta S(A \rightarrow C) + \Delta S(C \rightarrow D) - \Delta S(B \rightarrow D)$   
 $= 50 + 30 - 20 = +60 \text{ e.u.}$

6. (c)

7. (d): Energy left unutilized =  $\frac{1560}{2} = 780 \text{ kJ}$

Amount of water perspired

$$= \frac{\text{Energy left unutilized}}{\text{Enthalpy of evaporation of water}} = \frac{780}{44} \text{ mol}$$

$$\text{Amount of water (in g)} = \frac{780 \times 18}{44} = 319 \text{ g}$$

8. (c): The specific heat is an intensive property which does not depend on the quantity and size of matter.

9. (a):  $\Delta G^\circ = -2.303 RT \log K$   
 $= -2.303 \times 8.314 \times 298 \log (1.3 \times 10^3)$   
 $= -17767.688 \text{ J} = -17.767 \text{ kJ}$   
 $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$   
 $-17.767 = -66.5 - 298 \times \Delta S^\circ$   
 $\Delta S^\circ = \frac{-66.5 + 17.767}{298} = -0.163 \text{ kJ}$

10. (c): Exothermic compounds are more stable than endothermic compounds. Greater is the amount of heat evolved in the formation of a compound, more will be its stability.

11. (b, c, d)      12. (a, b, d)

13. (a, c, d):  $\Delta S_{\text{system}} = \Delta S_{\text{total}} - \Delta S_{\text{surrounding}}$

14. (a): At point (1), pressure is 1 atm and volume is 22.4 L for one mole of gas which indicates NTP state. Thus, temperature is 273 K. At point (2), pressure is again 1 atm but volume is doubled, so temperature is also doubled *i.e.*, 546 K. At point (3), pressure is halved but volume is doubled, so temperature is 273 K.

15. (b):  $\Delta E = nC_v\Delta T = 1 \times 3/2 \times 8.314 \times 273 = 3.4 \times 10^3 \text{ J}$

16. (a)      17. (d)      18. (b)

19. (b): (P):  $\Delta n_g = 2 - 2 = 0$ ; hence,  $\Delta H = \Delta U$   
 (Q):  $\Delta n_g = 2 - 1 = 1$ ; hence,  $\Delta H = \Delta U + RT$   
 (R):  $\Delta n_g = 2 - 4 = -2$ ; hence,  $\Delta H = \Delta U - 2RT$   
 (S):  $\Delta n_g = 5 - 2 = 3$ ; hence,  $\Delta H = \Delta U + 3RT$

20. (b)      21. (a)

22. (c): Spontaneity of a process depends upon  $\Delta G^\circ$  which combines with both factors,  $\Delta H^\circ$  and  $\Delta S^\circ$ .

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

For the spontaneity,  $\Delta G^\circ$  should be negative. If  $\Delta H^\circ$  is +ve,  $\Delta S^\circ$  is +ve then,  $\Delta G^\circ$  will be -ve only when  $T\Delta S > \Delta H$  which is possible at high temperature.

23. (4):  $\Delta S = \frac{q_{\text{rev}}}{T} = \frac{4520}{(-160 + 273)} = \frac{4520}{113} = 40 \text{ J mol}^{-1}$   
 $\Delta S = 10y = 40 \Rightarrow y = 4$

24. (0): At equilibrium,  $\Delta G = 0$

25. (9):  $C_v = 2.5 \text{ kJ K}^{-1} = 2500 \text{ J K}^{-1}$   
 $\Delta T = T_2 - T_1 = 298.45 - 298 = 0.45 \text{ K}$   
 $\Delta H$ , due to combustion of 3.5 g gas  
 $= C_v \times \Delta T = 2500 \times 0.45 = 1125 \text{ J}$   
 Given, molecular weight of gas = 28 = 1 mol  
 Hence,  $\Delta H$  due to combustion of 1 mole of gas  
 $= \frac{1125}{3.5} \times 28 = 9000 \text{ J mol}^{-1} = 9 \text{ kJ mol}^{-1}$

### Your favourite MTG Books/Magazines available in TRIPURA at

- Books Corner - Agartala Ph: 1381-2301945, 2301945; Mob: 9856358594
- Babai Books - Agartala Mob: 9774424611
- Puthi Ghar - Agartala Mob: 9436126357

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call

**0124-6601200** for further assistance.



YOUR WAY **CBSE XI**



Series 5

**CHAPTERWISE PRACTICE PAPER : HYDROGEN | THE s-BLOCK ELEMENTS**

Time Allowed : 3 hours

Maximum Marks : 70

**GENERAL INSTRUCTIONS**

- (i) All questions are compulsory.
- (ii) Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- (iii) Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- (iv) Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- (v) Q. no. 23 is a value based question and carries 4 marks.
- (vi) Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- (vii) Use log tables if necessary, use of calculators is not allowed.

1. Why are metallic hydrides also called interstitial hydrides?
2. Why is  $\text{Li}_2\text{CO}_3$  decomposed at a lower temperature?
3. What is meant by 15 volume of  $\text{H}_2\text{O}_2$  solution?
4. Why crystalline salts of alkaline earth metals contain more number of water of crystallisation than that of corresponding alkali metals?
5. Why  $\text{H}_2\text{O}_2$  cannot be stored for prolonged period?
6. Among  $\text{LiI}$  or  $\text{KI}$ , which one is more soluble in ethanol and why?
7. Answer the following questions :
  - (i) How would you prepare very pure dihydrogen in the laboratory?
  - (ii) How would you prepare heavy hydrogen in the laboratory?
8. Why is the temperature maintained around 393 K during the preparation of plaster of Paris?

**OR**

On exposure to air, sodium hydroxide becomes liquid and after sometimes it changes to white powder. Explain why?
9. Calculate the volume strength of a solution containing 30.36 g/lit of  $\text{H}_2\text{O}_2$ .
10. A compound X shows the following reactions :
  - (i) Zinc powder when boiled with a concentrated solution of X, it gets dissolved and hydrogen is evolved.
  - (ii) When an aqueous solution of X is added to an aqueous solution of stannous chloride, a white precipitate is obtained first which dissolves in excess of solution X.Identify the compound X and write down the reactions involved.
11. Arrange the following :
  - (i)  $\text{CaH}_2$ ,  $\text{BeH}_2$  and  $\text{TiH}_2$  in order of increasing electrical conductance.
  - (ii)  $\text{LiH}$ ,  $\text{NaH}$  and  $\text{CsH}$  in order of increasing ionic character.
  - (iii)  $\text{H}-\text{H}$ ,  $\text{D}-\text{D}$  and  $\text{F}-\text{F}$  in order of increasing bond dissociation enthalpy.
12. What happens when
  - (i) Magnesium is burnt in air
  - (ii) Quicklime is heated with silica
  - (iii) Calcium nitrate is heated?

13. Comment on the reactions of dihydrogen with  
 (i) Chlorine (ii) Sodium  
 (iii) Copper (II) oxide.
14. List three properties which illustrate the diagonal relationship between lithium and magnesium and three properties which illustrate the difference of lithium with other alkali metals.
15. How is hydrogen peroxide obtained in the laboratory? Write any two methods.
16. Explain the following :  
 (i) NaCl is not hygroscopic but table salt on exposure to air damps up.  
 (ii) Unlike  $\text{Na}_2\text{CO}_3$ ,  $\text{K}_2\text{CO}_3$  cannot be prepared by Solvay process.  
 (iii) It is necessary to add gypsum in the final stages of preparation of cement.
- OR**
- (i) Write down the reaction sequence to obtain :  
 (a) Bleaching powder from  $\text{CaCO}_3$ .  
 (b) Anhydrous  $\text{MgCl}_2$  from hydrated magnesium chloride.  
 (ii) Complete the following reaction :  
 $\text{Mg}(\text{HCO}_3)_2 + \text{Ca}(\text{OH})_2 \longrightarrow$
17. What is meant by water gas shift reaction? Discuss its use for the preparation of hydrogen.
18. Starting with sodium chloride how would you proceed to prepare (state the steps only)  
 (i) Sodium metal (ii) Sodium hydroxide  
 (iii) Sodium peroxide?
19. What do you understand by  
 (i) electron-deficient (ii) electron-precise and  
 (iii) electron-rich hydrides?  
 Provide justification with suitable examples.
20. Answer the following questions :  
 (i) Why can caesium be used in photoelectric cell while lithium cannot?  
 (ii) Lithium has highest ionization enthalpy in group I elements, yet it is the strongest reducing agent, why?  
 (iii) Why is calcium preferred over sodium to remove last traces of moisture from alcohol?
21. Calculate the normality of 20 volume solution of  $\text{H}_2\text{O}_2$ .
22. Compare the thermal stability of the following compounds of the alkali metals with those of alkaline earth metals.  
 (i) Nitrates (ii) Carbonates

23. Teacher told the students that water should be softened before using in boilers. Karina one of the students, asked the reason. The teacher explained the reason in detail to her and then Karina got satisfied.  
 (i) What explanation should be given by teacher?  
 (ii) What is temporary hardness?  
 (iii) Temporary hard water becomes soft on boiling. Why?  
 (iv) What values are shown by Karina?
24. Give the names and formula of the compounds indicated in the following statements:  
 (i) A compound of Ca used in setting fractured bones.  
 (ii) A compound of Mg, S, O and H used as a purgative in medicine.  
 (iii) A compound of Ca and C used for the production of acetylene (Give reaction also).  
 (iv) A compound of Ca, C and N used as a fertilizer.  
 (v) A triatomic compound of calcium which on treatment with water gives  $\text{H}_2$  (Give reaction also).

**OR**

Aqueous solution of an inorganic compound (X) shows the following reactions :

- (i) It decolourises an acidified  $\text{KMnO}_4$  solution accompanied by the evolution of oxygen.  
 (ii) It liberates iodine from an acidified KI solution.  
 (iii) It gives a brown precipitate with alkaline  $\text{KMnO}_4$  solution with evolution of oxygen.  
 (iv) It removes black stains from old oil paintings.  
 Identify (X) and give chemical equation for the reactions involved.
25. Complete the following chemical reactions.  
 (i)  $\text{Na}_{(s)} + \text{H}_{2(g)} \rightarrow$  (ii)  $\text{MnO}_4^-(\text{aq}) + \text{H}_2\text{O}_{2(\text{aq})} \rightarrow$   
 (iii)  $\text{CaO}_{(s)} + \text{H}_2\text{O}_{(g)} \rightarrow$  (iv)  $\text{AlCl}_{3(g)} + \text{H}_2\text{O}_{(l)} \rightarrow$   
 (v)  $\text{Ca}_3\text{N}_{2(s)} + \text{H}_2\text{O}_{(l)} \rightarrow$   
 Classify the above reactions into (a) hydrolysis (b) redox and (c) hydration.

**OR**

What do you understand by the terms :

- (i) hydrogen economy (ii) hydrogenation  
 (iii) syngas (iv) water-gas shift reaction  
 (v) hard water?
26. Explain the following :  
 (i) A sodium fire in laboratory is not extinguished by water.  
 (ii) Sodium and potassium do not form complex ions.



- (iii) Alkali metals are paramagnetic but their salts are diamagnetic.
- (iv) The softness of group IA metals increases down the group with increasing atomic number.
- (v) Alkali metals are obtained by the electrolysis of the molten salts and not by the electrolysis of their aqueous solutions.

**OR**


Explain the following :

- (i) Alkaline earth metals are weaker reducing agents than alkali metals.
- (ii) Beryllium shows covalency whereas other alkaline earth metals show electrovalency.
- (iii) Alkaline earth metals prefer to form  $M^{2+}$  ions.
- (iv) Be and Mg do not give colour to the flame whereas other alkaline earth metals do so.
- (v) The atomic radius of Mg is smaller than sodium.


### SOLUTIONS

1. In metallic hydrides, hydrogen is present in the interstices (holes or voids) existing in between the atoms in the lattice without changing the original crystal structure of the metal. That is why, these hydrides are called interstitial hydrides.
2. The  $\text{Li}^+$  ion exerts a strong polarising action and distorts the electron cloud of the nearby oxygen atom of the large  $\text{CO}_3^{2-}$  ion. This results in the weakening of the C—O bond and strengthening of the Li—O bond. Hence, it is less stable and decompose at lower temperature.
3. 15 volume  $\text{H}_2\text{O}_2$  solution means that 1 litre of this solution gives 15 litre of oxygen at S.T.P.
4. This is because of higher tendency of alkaline earth metal ions towards hydration as they have smaller ionic size and higher charge than alkali metals.
5.  $\text{H}_2\text{O}_2$  is an unstable compound which decomposes on standing to give water and oxygen, thus cannot be stored for a long period of time.
6.  $\text{LiI}$  is more soluble than  $\text{KI}$  in ethanol. Due to its small size, the lithium ion has a higher polarising power than potassium ion. It polarises the electron cloud of the iodide ion to a much greater extent than potassium ion. This causes a greater covalent character in  $\text{LiI}$  than  $\text{KI}$ . Hence,  $\text{LiI}$  is more soluble than  $\text{KI}$  in ethanol.
7. (i) Fairly pure hydrogen can be obtained by treating pure magnesium or pure aluminium with chemically pure  $\text{H}_2\text{SO}_4$  or  $\text{HCl}$  diluted with


distilled water. The gas is passed over  $\text{P}_2\text{O}_5$  and is collected by the displacement of mercury.



- (ii) Heavy hydrogen can be produced by the action of sodium metal on heavy water.



8. Plaster of Paris is prepared by heating gypsum to 393 K.




If the temperature is raised above 393 K, plaster of Paris is further dehydrated to form anhydrous calcium sulphate which is known as dead burnt plaster since it loses the property of setting with water.

**OR**

On exposure to air, sodium hydroxide continuously absorbs  $\text{CO}_2$  from atmosphere and converted into sodium carbonate. A stage reaches when the solution becomes saturated and crystals are formed. After some time, these  $\text{Na}_2\text{CO}_3$  crystals lose water of crystallisation (efflorescence) on standing in air and crumble in to white powder.

9. Volume strength of 30.36 g/lit  $\text{H}_2\text{O}_2$  means the volume of  $\text{O}_2$  (in L) that is liberated from 30.36 g of  $\text{H}_2\text{O}_2$ .






68 g of  $\text{H}_2\text{O}_2$  produce  $\text{O}_2$  at S.T.P. = 22.4 L

$$30.36 \text{ g of } \text{H}_2\text{O}_2 \text{ will produce } \text{O}_2 \text{ at S.T.P.} \\ = \frac{30.36 \times 22.4}{68} = 10 \text{ L}$$

$\therefore$  Volume strength of  $\text{H}_2\text{O}_2$  = 10 L.

10. The compound X is  $\text{NaOH}$  and the reactions involved are



11. (i)  $\text{BeH}_2$  is significantly covalent,  $\text{CaH}_2$  is ionic and  $\text{TiH}_2$  is metallic hydride. Hence, increasing electrical conductance :




(ii) Electronegativity decreases as  $\text{Li} > \text{Na} > \text{Cs}$ .



Thus, increasing ionic character :




(iii) Due to lone pairs of F, bond pairs experience repulsion, hence, F-F has low bond dissociation energy. In D-D, due to higher nuclear attraction bond dissociation energy is greater than H-H. Increasing bond dissociation enthalpy :




12. (i) When magnesium is burnt in air it forms magnesium oxide and magnesium nitride with dazzling brilliance.




(ii) When quicklime is heated with silica, it gives calcium silicate.




(iii) When calcium nitrate is heated, it gives  $\text{CaO}$ ,  $\text{NO}_2$  and  $\text{O}_2$ .




13. (i) Dihydrogen reduces chlorine to chloride ion ( $\text{Cl}^-$ ) and itself gets oxidised to  $\text{H}^+$  ion in  $\text{HCl}$ . An electron pair is shared between H and Cl to form a covalent molecule of hydrogen chloride.



(ii) Sodium reduces dihydrogen to form hydride ( $\text{H}^-$ ) ion and itself gets oxidised to sodium ( $\text{Na}^+$ ) ion. During this reaction, an electron is transferred from Na to H leading to the formation of an ionic compound of sodium hydride ( $\text{NaH}$ ).





(iii) Hydrogen reduces copper (II) oxide to copper metal (in zero oxidation state) and itself gets oxidised to  $\text{H}_2\text{O}$ , which is a covalent molecule.



14. Properties which illustrate the diagonal relationship between Li and Mg :

(i) Both Li and Mg are harder and have higher melting points than other elements in their respective groups.

(ii) Lithium and magnesium both form only monoxides when heated in oxygen.





(iii) Chlorides of both Li and Mg are deliquescent and soluble in alcohol and pyridine.

Properties which illustrate the point of difference of Li to other alkali metals :

(i) Li is much harder. Its m.pt. and b.pt. are higher than the other alkali metals.


(ii) Li is least reactive but the strongest reducing agent among all the other alkali metals.

(iii)  $\text{LiNO}_3$ , on heating gives a mixture of  $\text{NO}_2$  and  $\text{O}_2$  while nitrates of rest of alkali metals gives only oxygen.




15. In laboratory, hydrogen peroxide may be prepared from sodium peroxide and barium peroxide.

(i) Sodium peroxide is added in small amount to an ice cold dilute solution (20 per cent) of sulphuric acid.



Sodium sulphate formed is removed by crystallisation, and a dilute solution of  $\text{H}_2\text{O}_2$  is obtained.


(ii) Paste of  $\text{BaO}_2 \cdot 8\text{H}_2\text{O}$  is prepared and added gradually to an ice cold dilute sulphuric acid. The solution is kept well-stirred throughout.



Barium sulphate precipitates out leaving behind a dilute solution of hydrogen peroxide.

16. (i) Pure  $\text{NaCl}$  is not hygroscopic. The commercial common salt or table salt consists of small amount of  $\text{MgCl}_2$  and  $\text{CaCl}_2$  which absorb moisture from atmosphere making it damp.

(ii) In Solvay process,  $\text{NaHCO}_3$  is prepared first which becomes sparingly soluble and on heating,  $\text{Na}_2\text{CO}_3$  is obtained.




$\text{KHCO}_3$  is soluble in water and can not be obtained in solid state that is why  $\text{K}_2\text{CO}_3$  is not formed by this method.


(iii) When cement is mixed with water, it becomes hard over a period of time. This is called setting of cement. Gypsum is often added to portland cement to prevent early hardening or flash setting allowing a longer working time. Gypsum slows down the setting of cement so that cement is adequately hardened.

OR

- (i) (a) Getting bleaching powder from  $\text{CaCO}_3$ .





- (b) Anhydrous  $\text{MgCl}_2$  from hydrated  $\text{MgCl}_2$  : To make hydrated  $\text{MgCl}_2$  anhydrous, it is first heated in air and then in a current of  $\text{HCl}$  gas.



- (ii)  $\text{Mg(HCO}_3)_2 + \text{Ca(OH)}_2 \longrightarrow \text{MgCO}_3 + \text{CaCO}_3 + 2\text{H}_2\text{O}$


17. Water gas shift reaction involves a selective oxidation of  $\text{CO}$  to  $\text{CO}_2$  by passing a mixture of water gas and steam (excess) over a catalyst containing ferric oxide and chromium oxide at 773 K. Water gas is obtained by passing steam over red hot coke.



$\text{CO}_{2(g)}$  is removed by passing the mixture through water under high pressure. Thus, water gas shift reaction generates more hydrogen.


18. (i)  $\text{NaCl}_{(s)} \xrightarrow{\text{heat}} \text{NaCl} \xrightarrow[\text{(Molten)}]{\text{electrolysis}} \text{Na}_{(s)} + \frac{1}{2}\text{Cl}_{2(g)}$   
at cathode at anode

- (ii)  $\text{NaCl}_{(aq)} \xrightarrow[\text{Hg-cathode}]{\text{electrolysis}} \text{Na} - \text{Hg} + \frac{1}{2}\text{Cl}_{2(g)}$   
at cathode at anode



- (iii)  $\text{NaCl}_{(s)} \xrightarrow{\text{heat}} \text{NaCl} \xrightarrow[\text{(molten)}]{\text{electrolysis}} \text{Na}_{(s)} + \frac{1}{2}\text{Cl}_{2(g)}$   
at cathode at anode

This sodium is then heated on aluminium trays in air (free of  $\text{CO}_2$ ) to form its peroxide.



19. (i) Electron-deficient hydrides are those which do not have sufficient number of electrons to form normal covalent bonds. For example, hydrides of group 13 ( $\text{BH}_3$ ,  $\text{AlH}_3$ , etc). To make up their

deficiency they generally exist in polymeric forms such as  $\text{B}_2\text{H}_6$ ,  $\text{Al}_2\text{H}_6$ , etc.

(ii) Electron-precise hydrides are those which have sufficient number of electrons required for forming covalent bonds. For example, hydrides of group 14 ( $\text{CH}_4$ ,  $\text{SiH}_4$ ,  $\text{GeH}_4$ ,  $\text{SnH}_4$ ,  $\text{PbH}_4$  etc).


(iii) Electron-rich hydrides are those which have excess electrons required to form normal covalent bonds. For example, hydride of group 15 or 17 (i.e.,  $\text{NH}_3$ ,  $\text{PH}_3$ ,  $\text{H}_2\text{O}$ ,  $\text{H}_2\text{S}$ ,  $\text{H}_2\text{Se}$ ,  $\text{H}_2\text{Te}$ , etc). The excess electrons in these hydrides are present as lone pairs.

20. (i) Caesium has the lowest while lithium has the highest ionization enthalpy. Hence, Cs can lose electrons very easily while lithium cannot and hence can not be used in photoelectric cell.

(ii) In aqueous solutions, the tendency of an element to lose electrons does not entirely depend upon its ionization enthalpy. It also depends upon its enthalpy of sublimation and the enthalpy of hydration of the ion left after the loss of an electron. The combined effect of these factors is measured in terms of electrode potential. Since lithium has the most negative electrode potential in group I elements, therefore, lithium is the strongest reducing agent.

(iii) Both Na and Ca react with water to form their respective hydroxides. On the other hand, Na readily reacts with alcohol to form sodium ethoxide but Ca reacts very slowly. Thus, when Ca is used, it will react more readily with water than with alcohol.  
 $2\text{C}_2\text{H}_5\text{OH} + 2\text{Na} \longrightarrow 2\text{C}_2\text{H}_5\text{ONa} + \text{H}_2$   
Therefore, Ca is preferred over Na to remove the last traces of moisture from alcohol.

21.  $\text{H}_2\text{O}_2$  decomposes as follows :



Let us take 1 L of 20 volume  $\text{H}_2\text{O}_2$ . Then,

1 L of  $\text{H}_2\text{O}_2 \equiv 20$  L of oxygen

From the stoichiometry,

22.4 L of oxygen is given by 4 equiv. of  $\text{H}_2\text{O}_2$


1 L of oxygen is given by  $\frac{4 \text{ equiv.}}{22.4}$  of  $\text{H}_2\text{O}_2$

20 L of oxygen is given by  $\frac{4 \text{ equiv.}}{22.4} \times 20 = 3.57$  equiv.


Thus, 1 L of  $\text{H}_2\text{O}_2$  contains 3.57 equivalents of  $\text{H}_2\text{O}_2$  in it.

$\therefore$  Normality of  $\text{H}_2\text{O}_2$  sample = 3.57


22. (i) Thermal stability of nitrates : Nitrates of both alkali and alkaline earth metals decompose on heating. All alkaline earth metal nitrates decompose to form metal oxide,  $\text{NO}_2$  and  $\text{O}_2$ .




The nitrates of Na, K, Rb and Cs decompose to form metal nitrites and  $\text{O}_2$ .



However, due to diagonal relationship between Li and Mg, lithium nitrate decomposes like  $\text{Mg}(\text{NO}_3)_2$  to form metal oxide,  $\text{NO}_2$  and  $\text{O}_2$ .




- (ii) Thermal stability of carbonates : Carbonates of alkaline earth metals decompose on heating to form metal oxide and  $\text{CO}_2$ .



Further as the electropositive character of the metal increases down the group, the stability of these metal carbonates increases or the temperature of their decomposition increases as shown below :

$\text{BeCO}_3$	$\text{MgCO}_3$	$\text{CaCO}_3$	$\text{SrCO}_3$	$\text{BaCO}_3$
373 K	813 K	1173 K	1563 K	1633 K

Due to diagonal relationship between Li and Mg,  $\text{Li}_2\text{CO}_3$  decomposes in the same way as  $\text{MgCO}_3$ .



All other alkali metal carbonates are stable and do not decompose even at high temperature.

23. (i) Water should be softened before using in boilers otherwise it may cause various boiler problems like



(a) Scale and sludge formation

(b) Priming and foaming

(c) Boiler corrosion.

(ii) Temporary hardness of water is the hardness caused by the presence of dissolved calcium and magnesium bicarbonates.

(iii) On boiling the calcium/magnesium bicarbonate decomposes to give calcium/magnesium carbonate, which is insoluble in water. Therefore, it precipitates out.




- (iv) Curiosity, Knowledge and self awareness.

24. (i) Plaster of Paris,  $2\text{CaSO}_4 \cdot \text{H}_2\text{O}$


(ii) Magnesium sulphate,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$

(iii) Calcium carbide,  $\text{CaC}_2$








(iv) Calcium cyanamide (Nitrolim),  $\text{CaCN}_2 + \text{C}$

(v) Calcium hydride (hydrolith),  $(\text{CaH}_2)$







**OR**



Therefore, compound (X) is  $\text{H}_2\text{O}_2$ .



25. (i)  $2\text{Na} + \text{H}_2 \rightarrow 2\text{NaH}$  (Redox reaction)



**OR**

(i) Hydrogen economy : The energy is transported and stored in the form of liquid or gaseous hydrogen which is considered to be a possible source of clean energy. This proposal to use hydrogen as fuel is called hydrogen economy.


(ii) Hydrogenation : The process of addition of hydrogen to unsaturated hydrocarbons is known as hydrogenation.



(iii) Syngas : The mixture of  $\text{CO}$  and  $\text{H}_2$  is used for the synthesis of methanol and a number of hydrocarbons, it is also called synthesis gas or syngas. Nowadays, syngas is produced from sewage, saw dust, scrap wood, newspapers, etc.



(iv) Water-gas shift reaction : Commercial dihydrogen is obtained by mixing water gas with steam and passed over a mixture of ferric oxide and chromium oxide heated at 500 °C.



This is called water-gas shift reaction.

(v) Hard water : The water containing calcium and magnesium salts in the form of hydrogen carbonate, chloride and sulphate is known as hard water.

26. (i) Since sodium reacts vigorously with water, sodium fire should not be extinguished with water, but carbon tetrachloride (pyrene) should be used to extinguish sodium fire in laboratory.

(ii) In order to form complex compounds, a metal should have the vacant *d*-orbitals which Na or K or other alkali metals do not have so cannot form complex ions.

(iii) The alkali metals have one unpaired electron in their outermost shell, thus these are paramagnetic. While in their salts, alkali metals are present as cations which have inert gas configuration thus their salt are diamagnetic.

(iv) The softness of group IA metals increases down the group because there is continuous decrease of metallic bond strength on account of an increase in atomic size. Bigger is the atomic size of metal, its metallic bond strength is weak and metal becomes soft.

(v) The aqueous solution of alkali metal salt contains metal cation,  $\text{H}^+$ ,  $\text{OH}^-$  and anion. In that condition the discharge potential of  $\text{H}^+$  ions is lower than the metal cations and on electrolysis of solutions of alkali metal salts, hydrogen is discharged at cathode rather than the metal.

OR

(i) Since the oxidation potential of alkaline earth metals is lower than that of alkali metals, thus alkaline earth metals are weaker reducing agents than alkali metals.

(ii) The polarising power of  $\text{Be}^{2+}$  ion is high, due to its high charge density. The large distortion of electron cloud of the anion brings covalent character. Be has high value of electronegativity and when it reacts with another element, the electronegativity difference is not much and the bond formed is therefore covalent.

(iii) The existence of divalent ions in solid state is due to greater lattice energy released in the formation of compounds of divalent cations than to those of monovalent cations. High value of hydration energy of divalent cations counter balances the higher value to ionisation energy. Thus, alkaline earth metals prefer to form  $M^{2+}$  ions.

(iv) In case of Ca, Sr, Ba, Ra, the electron can be excited by the supply of energy. When excited electron comes in its original position it releases energy in the form of light. But Be and Mg have small size, thus their electrons are tightly held and thus excitation of electron is difficult. Hence, they do not show flame colouration.

(v) Shell number of both Na and Mg is same. But number of electrons in Mg is more, thus attractive force in Mg is greater than that in Na and thus Mg has smaller size than Na.

### Your favourite MTG Books/Magazines available in PUNJAB at

- Sunder Book Depot - Amritsar  
Ph: 0183-2544491, 2224884, 5059033; Mob: 9814074241
- Malhotra Book Depot - Amritsar Mob: 8000300086, 9646537157, 9888733844
- Navchattan Book Depot - Barnala  
Mob: 97790220692, 9417906880, 9779090135, 9876147263, 9779050692
- Mehta Book Centre - Bathinda Mob: 9876029048, 9464158497
- Goyal Traders Book Sellers - Bathinda  
Ph: 0164-2239632; Mob: 9417924911, 9814485520
- Aggarwal Book Centre - Bathinda Ph: 0164-2236042; Mob: 9417816439
- S M Enterprises - Bathinda Ph: 0164-2240450; Mob: 7508622881, 9417363362
- Hans Raj Dogra & Sons - Gurdaspur Mob: 9872032357
- Gupta Technical & Computer Book Shop - Jalandhar  
Ph: 0181-2200397; Mob: 9915001916, 9779981081
- Cheap Book Store - Jalandhar Mob: 9872223458
- City Book Shop - Jalandhar Ph: 0181-2620800; Mob: 9417440753
- Deepak Book Distributions - Jalandhar  
Ph: 0181-2222131; Mob: 8528391133, 9872328131
- Kiran Book Shop - Jalandhar Mob: 9876631526, 9779223883, 9872377808
- Amit Book Depot - Ludhiana  
Ph: 0161-5022930, 5022930; Mob: 9815323429, 9815807871
- Bhatia Book Centre - Ludhiana Ph: 0161-2747713; Mob: 9815277131
- Chabra Book Depot - Ludhiana Ph: 0161-6900900, 2405427; Mob: 9501080070
- Gupta Book World - Ludhiana  
Ph: 0161-2446870, 2409097, 3942992; Mob: 9463027555
- Khanna Book Depot - Nabha Ph: 01765-220095; Mob: 9814093193, 9814309320
- Goel & Sons Book Depot - Patiala Ph: 0172-2213643, 2202045; Mob: 9914096991
- Adarsh Pustak Bhandar - Patiala  
Ph: 0175-2311430; Mob: 9814347613, 9815651737
- Pepsu Book Depot - Patiala Ph: 0175-2214696; Mob: 9814041623, 9914903305

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call **0124-6601200** for further assistance.

# MPP-7 MONTHLY Practice Problems

Class XI

This specially designed column enables students to self analyse their extent of understanding of specified chapter. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.



## The s-Block Elements

Total Marks : 120

Time Taken : 60 Min.

NEET / AIIMS

Only One Option Correct Type

- When  $\text{NaNO}_2$  is treated with Na
  - $\text{NaNO}_3$  is formed
  - $\text{Na}_2\text{O}$  and  $\text{N}_2$  are formed
  - $\text{Na}_2\text{O}_2$  and  $\text{N}_2$  are formed
  - $\text{Na}_2\text{O}$  and  $\text{Na}_3\text{N}$  are formed.
- Sodium metal reacts with  $\text{Al}_2\text{O}_3$  at high temperature to give a sodium compound X. X reacts with carbon dioxide in water to form Y. Y is
  - $\text{Na}_2\text{CO}_3$
  - $\text{Na}_2\text{O}_2$
  - $\text{NaAlO}_2$
  - $\text{Na}_2\text{O}$
- In Solvay process,  $\text{NaCl}$  combines with  $\text{NH}_4\text{HCO}_3$  to give
  - precipitate of  $\text{NaHCO}_3$
  - precipitate of  $\text{Na}_2\text{CO}_3$
  - solution of  $\text{NaHCO}_3$
  - solution of  $\text{Na}_2\text{CO}_3$ .
- Elements of group I give colour to the Bunsen burner flame due to
  - softness
  - low melting point
  - low ionisation enthalpy
  - single electron in the outermost orbit.
- Stability of nitrides of IIA group elements is in order
  - $\text{Ca}_3\text{N}_2 < \text{Mg}_3\text{N}_2 < \text{Be}_3\text{N}_2$
  - $\text{Be}_3\text{N}_2 < \text{Mg}_3\text{N}_2 < \text{Ca}_3\text{N}_2$
  - $\text{Ca}_3\text{N}_2 < \text{Be}_3\text{N}_2 < \text{Mg}_3\text{N}_2$
  - $\text{Mg}_3\text{N}_2 < \text{Be}_3\text{N}_2 < \text{Ca}_3\text{N}_2$
- Consider the following abbreviations for hydrated alkali ions
 
$$X = [\text{Li}(\text{H}_2\text{O})_n]^+ \quad Y = [\text{K}(\text{H}_2\text{O})_n]^+ \\ Z = [\text{Cs}(\text{H}_2\text{O})_n]^+$$
 Which of the following is the correct order of size of these hydrated alkali ions?
  - $Z > X > Y$
  - $X = Y = Z$
  - $X > Y > Z$
  - $Z > Y > X$
- Calcium is obtained by
  - electrolysis of molten  $\text{CaCl}_2$
  - electrolysis of solution of  $\text{CaCl}_2$  in water
  - reduction of  $\text{CaCl}_2$  with carbon
  - roasting of limestone.
- Which of the following is not correct about the solution when moderate amount of sodium metal is dissolved in liquid ammonia at low temperature?
  - $\text{Na}^+$  ions are produced in solution.
  - A blue coloured solution is obtained.
  - The solution acts as a good conductor of electricity.
  - Liquid ammonia remains diamagnetic.
- What happens when quick lime is exposed to air?
  - It produces  $\text{CaCO}_3$  and  $\text{H}_2\text{O}$ .
  - It produces  $\text{Ca}(\text{OH})_2$  and  $\text{H}_2\text{O}$ .
  - It produces  $\text{Ca}(\text{OH})_2$  and  $\text{CaCO}_3$ .
  - It produces  $\text{CaH}_2$  and  $\text{H}_2\text{O}$ .
- By adding gypsum to cement
  - setting time of cement becomes less
  - setting time of cement increases
  - colour of cement becomes light
  - shining surface is obtained.

11. Comparing calcium oxide and magnesium oxide, which of the two would be more stable?

- Magnesium oxide would be more stable because the magnesium cation is smaller than the calcium cation.
- Magnesium oxide would be more stable because magnesium has lower atomic mass than calcium.
- Calcium oxide would be more stable because the calcium cation is smaller than the magnesium cation.
- Calcium oxide would be more stable because calcium has greater atomic mass than magnesium.

12. The pair of compounds which cannot exist together in solution is

- $\text{NaHCO}_3$  and  $\text{NaOH}$
- $\text{Na}_2\text{CO}_3$  and  $\text{NaHCO}_3$
- $\text{Na}_2\text{CO}_3$  and  $\text{NaOH}$
- $\text{NaHCO}_3$  and  $\text{NaCl}$ .

#### Assertion & Reason Type

**Directions :** In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- If both assertion and reason are true and reason is the correct explanation of assertion.
- If both assertion and reason are true but reason is not the correct explanation of assertion.
- If assertion is true but reason is false.
- If both assertion and reason are false.

13. **Assertion :** Among the alkali metals, lithium salts exhibit the least electrical conductance in aqueous solutions.

**Reason :** Smaller the radius of the hydrated cation, lower is the electrical conductance in aqueous solutions.

14. **Assertion :**  $\text{Na}_2\text{SO}_4$  is soluble in water but  $\text{BaSO}_4$  is insoluble in water.

**Reason :** In case of  $\text{BaSO}_4$ , its lattice energy is more than its hydration energy.

15. **Assertion :** Barium is not required for normal biological functions in human.

**Reason :** Barium does not show variable oxidation states.

#### JEE MAIN / JEE ADVANCED

##### Only One Option Correct Type

16.  $\text{Na}_2\text{O}_2$

- is diamagnetic in nature
- is a salt of dibasic acid  $\text{H}_2\text{O}_2$
- oxidises  $\text{Cr}^{3+}$  (green) to  $\text{CrO}_4^{2-}$  (yellow)
- all of these.

17. All the products are same when X and Y react with water. X and Y are respectively :

- Ca,  $\text{CaO}$
- Na,  $\text{NaH}$
- $\text{NaOH}$ ,  $\text{Na}_2\text{CO}_3$
- Na, K

18. A metal M readily forms water soluble sulphate  $\text{MSO}_4$ , water insoluble hydroxide  $\text{M}(\text{OH})_2$  and oxide MO which becomes inert on heating. The hydroxide is soluble in  $\text{NaOH}$ . Thus, M is

- Mg
- Sr
- Be
- Ca

19. The alkali metals form hydrides by direct synthesis at elevated temperature. The thermal stability of these hydrides decreases in which of the following order?

- $\text{CsH} > \text{RbH} > \text{KH} > \text{NaH} > \text{LiH}$
- $\text{KH} > \text{NaH} > \text{LiH} > \text{CsH} > \text{RbH}$
- $\text{NaH} > \text{LiH} > \text{KH} > \text{RbH} > \text{CsH}$
- $\text{LiH} > \text{NaH} > \text{KH} > \text{RbH} > \text{CsH}$

##### More than One Options Correct Type

20. Select the correct statements.

- Alkali metals form alloys amongst themselves.
- Alkali metals dissolve readily in mercury.
- The process of formation of amalgams by alkali metals is highly exothermic.
- The process of formation of amalgam by alkali metals is highly endothermic.

21. Which of the following compounds are used for textile industry?

- $\text{Na}_2\text{CO}_3$
- $\text{NaHCO}_3$
- $\text{NaOH}$
- $\text{NaCl}$

22. The compounds formed upon combustion of sodium metal in excess air are


- $\text{Na}_2\text{O}_2$
- $\text{Na}_2\text{O}$
- $\text{NaO}_2$
- $\text{NaOH}$

23. In the diaphragm cell used for the electrolysis of brine, the reactions that occur are

- $2\text{Na} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2$
- $2\text{NaOH} + \text{Cl}_2 \rightarrow \text{NaCl} + \text{NaOCl} + \text{H}_2\text{O}$
- $4\text{OH}^- \rightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^-$
- $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2\text{e}^-$

**Integer Answer Type**

24. Out of Be, Mg, Ca, Sr and Ba, total number of metals whose nitrates are decomposed according to following reaction :



25. Total number of alkali metals (non-radioactive) which produce stable superoxide when burn with excess of  $O_2$  is
26. Total number of moles of water in 1 mole of epsom salt is

**Comprehension Type**

On treatment with cold water, element (A) reacted quietly, liberating colourless gas (B) and solution (C). Lithium reacted with gas (B) giving a solid product (D) which gave effervescence with water to give basic solution (F). When  $CO_2$  was passed through solution (C), initially white ppt. (E) was formed but this redissolved forming solution (G) when more  $CO_2$  was passed. Ppt. (E) effervescenced when moistened with conc. HCl and gave a deep red colouration to Bunsen flame. (E) on heating with excess of carbon at  $2000^\circ C$  gave (H).

27. Consider the following statements about a salt solution (G).
- (I) It causes permanent hardness of water.
  - (II) It cannot be obtained in solid state.
  - (III) It causes temporary hardness to water.
  - (IV) It can be obtained in solid state.
- Correct statements are
- (a) I and II only
  - (b) I and IV only
  - (c) II and III only
  - (d) II and IV only
28. Solid (H) on hydrolysis gives a gas, which on passing through ammonical  $AgNO_3$ , gives
- (a) white ppt.
  - (b) red ppt.
  - (c) no ppt.
  - (d) brown ppt.

**Matrix Match Type**

29. Match the Column I with Column II and choose the correct option using the codes given below.

Column I (Compound)	Column II (Use)
(A) $CaCO_3$	(p) Used in gun powder
(B) $Na_2S_2O_3 \cdot 5H_2O$	(q) Used in fire extinguishers
(C) $NaNO_3$	(r) Used as a flux in metallurgy
(D) $NaHCO_3$	(s) Used in black and white photography

A	B	C	D
(a) r	s	p	q
(b) p	s	q	r
(c) r	q	p	s
(d) r	s	q	p

30. Match the Column I with Column II and choose the correct option using the codes given below.

Column I	Column II
(A) Complex formation	(p) Be
(B) Formation of covalent compounds	(q) Mg
(C) High solubility of salts	(r) Ca
(D) Explosive reaction with acids	(s) Sr

A	B	C	D
(a) q	p	p, q	q, s
(b) p	p	p, q	r, s
(c) p, s	q	r	s
(d) p	q	r, s	p, q



Keys are published in this issue. Search now! ☺

**SELF CHECK****Check your score! If your score is**

No. of questions attempted .....  
 No. of questions correct .....  
 Marks scored in percentage .....

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.



# NEET | JEE ESSENTIALS

Class  
XII

Maximise your chance of success, and high rank in NEET, JEE (Main and Advanced) by reading this column. This specially designed column is updated year after year by a panel of highly qualified teaching experts well-tuned to the requirements of these Entrance Tests.

## Unit 5

### Haloalkanes and Haloarenes | Alcohols, Phenols and Ethers

#### HALOALKANES AND HALOARENES

#### HALOALKANES

##### INTRODUCTION

- Compounds derived from alkanes by the replacement of one or more hydrogen atoms by the corresponding number of halogen atoms are termed as halogen derivatives of alkanes or haloalkanes.

HALOALKANES

**Monohalogen derivatives :**  $C_nH_{2n+2} \xrightarrow[-X]{-H} C_nH_{2n+1}X$ 

$CH_3Cl$   
Methyl chloride

$C_2H_5Br$   
Ethyl bromide

**Di-halogen derivatives :**  $C_nH_{2n+2} \xrightarrow[+2X]{-2H} C_nH_{2n}X_2$ 

<b>Gem-dihalide</b> Both halogen atoms are attached to same carbon atom. <i>e.g.</i> , $CH_3CH_2CHCl_2$ 1, 1-dichloropropane	<b>Vic-dihalide</b> The halogen atoms are attached to adjacent carbon atom, known as alkylene halides. <i>e.g.</i> , $ClCH_2-CH_2Cl$ 1, 2-dichloroethane	<b><math>\alpha-\omega</math> halides</b> The halogen atoms are attached to terminal carbon atoms. $\begin{array}{c} CH_2-CH_2-CH_2 \\   \qquad \qquad   \\ Br \qquad \qquad Br \end{array}$ 1, 3-dibromopropane
---	---	--

**Tri-halogen derivatives :**  $C_nH_{2n+2} \xrightarrow[+3X]{-3H} C_nH_{2n-1}X_3$ 

$CHCl_3$   
Chloroform


$CHBr_3$   
Bromoform

$CHI_3$   
Iodoform

**Tetrahalogen derivatives :** Four hydrogen atoms are replaced by four halogen atoms. *e.g.*,  $CCl_4$ ,  $CBr_4$ ,  $Cl_2HC-CHCl_2$ , etc.


## PREPARATION

- From alcohols :



Most preferred method is (III) because other products of this reaction ( $\text{SO}_2$  and  $\text{HCl}$ ) are gaseous and can easily escape.


- From hydrocarbons :




- Free radical mechanism is followed.
- Order of ease of substitution :  $3^\circ > 2^\circ > 1^\circ$  (alkyl)  
Benzylic  $\approx$  allylic  $>$  alkyl  $>$  vinylic  $\approx$  aryl

- From alkene (by HX addition) :

- From symmetrical alkenes :



- From unsymmetrical alkenes :




- Markovnikov's rule :** During the addition across unsymmetrical double bond, the negative part of the attacking reagent goes to the carbon atom carrying lesser number of hydrogen atoms while the positive part goes to the carbon atom with more number of hydrogen atoms.
- Anti-Markovnikov's rule :** When HBr is added to unsymmetrical double bond in presence of peroxide, the reaction takes place contrary to the Markovnikov's rule. This is known as anti-Markovnikov's rule, peroxide effect or Kharasch

effect. This law is applicable for the addition of HBr only.


- From silver salts of acids :

- Borodine Hunsdiecker reaction :




- By halogen exchange method :

- Finkelstein reaction



- Swartz reaction (For alkyl fluoride)




## PHYSICAL PROPERTIES

- Alkyl halides are very slightly soluble in water but soluble in organic solvents.
- Alkyl bromides and iodides are heavier than water whereas alkyl chlorides and fluorides are lighter than water.
- The boiling points of alkyl halides are in the order  $\text{RI} > \text{RBr} > \text{RCl} > \text{RF}$  and the boiling points of alkyl halides increases with increase in size of the alkyl group.
- Alkyl halides are in general toxic compounds and bring unconsciousness when inhaled in large amount.


## NATURE OF C—X BOND

- In haloalkanes, the carbon is bonded to a halogen atom ( $X = \text{F, Cl, Br, I}$ ) which is more electronegative than carbon. Consequently, the C—X bond is polar in nature and its bond length increases from C—F to C—I.



## CHEMICAL PROPERTIES

- Nucleophilic substitution reactions :** The reactions in which a stronger nucleophile displaces a weaker nucleophile (leaving group) are called nucleophilic substitution reactions.




- Leaving group ability :  $\text{I}^- > \text{Br}^- > \text{Cl}^- > \text{F}^-$

- Order of reactivity :  $\text{R}-\text{I} > \text{R}-\text{Br} > \text{R}-\text{Cl} > \text{R}-\text{F}$


## Types of Nucleophilic Substitution Reactions

$S_N2$ (Bimolecular)	$S_N1$ (Unimolecular)
<p><b>Mechanism :</b></p> <p>Rate = <math>k[RX][OH^-]</math></p> <p>Single step reaction.</p> <p>Inversion of configuration takes place.</p> <p>Reactivity order of alkyl halides : <math>CH_3X &gt; 1^\circ &gt; 2^\circ &gt; 3^\circ</math></p>	<p><b>Mechanism :</b></p> <p><b>Step I :</b> <math>CH_3-C(CH_3)_2-Br \rightleftharpoons CH_3-C^+(CH_3)_2 + Br^-</math></p> <p><b>Step II :</b> <math>CH_3-C^+(CH_3)_2 + OH^- \longrightarrow CH_3-C(CH_3)_2-OH</math></p> <p>Rate = <math>k[RX]</math></p> <p>Two step reaction.</p> <p>Carbocation rearrangement may take place.</p> <p>Reactivity order of alkyl halides : <math>3^\circ &gt; 2^\circ &gt; 1^\circ &gt; CH_3X</math></p>


### Examples of substitution reactions :



- Optical Activity :** Compounds which rotate the plane of polarised light when they pass through the prism, are called optically active compounds and the phenomenon is known as optical activity.




- Inversion, retention and racemisation :** When a reaction takes place at an asymmetric carbon atom following reactions are possible :




- Elimination reactions :**

- **Dehydrohalogenation :** These reactions are called β-elimination reactions which follow Saytzeff's rule.




- **Saytzeff rule :** In haloalkane, when elimination of hydrogen halide in two different ways occur, then that alkene will be preferred, in which carbon atoms joined by the double bond are maximum alkylated *i.e.*, contain largest number of alkyl groups.


- **Ease of formation of alkene :**



- **Ease of dehydrohalogenation :**





- Reactions with metals :**




## HALOARENES

### PREPARATION



- Hunsdiecker reaction :**




## PHYSICAL PROPERTIES

- Aryl halides are colourless stable liquids with pleasant odour, insoluble in water but readily miscible with organic solvents.
- Their boiling points are higher than corresponding alkyl halides. The boiling points rise gradually from fluoro to iodo compounds.

## CHEMICAL PROPERTIES

- Nucleophilic substitution reactions :** Haloarenes are less reactive towards nucleophilic substitution reactions:

- Due to resonance effect :



Thus, C—Cl bond acquires double bond character and difficult to break.


- C atom in haloarenes is  $sp^2$  hybridised, thus has greater  $s$ -character, so difficult to break.
- Instability of phenyl cation : Phenyl cation formed by self-ionisation will not be stabilised hence,  $S_N1$  mechanism ruled out.

### MPP-7 CLASS XI


### ANSWER KEY

- (b)
- (a)
- (a)
- (d)
- (a)
- (c)
- (a)
- (d)
- (c)
- (b)
- (a)
- (c)
- (a)
- (b)
- (d)
- (b)
- (c)
- (d)
- (a,b,c)
- (a,c)
- (a,b)
- (a,b,c,d)
- (5)
- (3)
- (7)
- (c)
- (a)
- (b)






- Electrophilic substitution reactions :** As electron density increases at *o*, *p*-positions, electrophilic substitution reactions occur at *o*, *p*-positions.



#### Other reactions :



## INFOSHOTS

### Halogen derivative of 1, 3-disubstituted thiourea!

1, 3-disubstituted thiourea derivatives have attracted an increasing attention due to their potential in medicinal chemistry. Among them, the broad spectrum of biological activities of fluorinated thiourea derivatives has been investigated and various bioactivities, such as antimicrobial, antiviral and anticancer have been reported. In recent years, several chlorophenyl, fluorophenyl and (trifluoromethyl) phenyl thiourea compounds have been described as potent antistaphylococcal and antifungal agents.

### Your favourite MTG Books/Magazines available in ORISSA at

- Padmalaya Book Seller - Bhubaneswar  
Ph: 0674-2396922, 3039692; Mob: 9437026922
- Pragnya - Bhubaneswar Ph: 0674-2395757, 30396922; Mob: 9861135495
- Sri Ram Books - Bhubaneswar Ph: 0674-2391385; Mob: 9439921093
- Sagar Book Store - Bhubaneswar Ph: 0674-2506040, 2516040; Mob: 8763406040
- Sri Madhab Book Store - Cuttack Ph: 0671-2431158, 2431148; Mob: 9437024244
- A.K. Mishra Agency Pvt. Ltd. - Cuttack  
Ph: 0671-2322244, 2322255, 2322266, 2332233; Mob: 9437025991
- Kitab Mahal - Cuttack Ph: 0671-2648333, 2648301; Mob: 9861388145

Visit **"MTG IN YOUR CITY"** on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call **0124-6601200** for further assistance.

# CONCEPT MAP

## CLASS XI


## Gaseous State

A gas occupies a volume about 1600 times greater than that of an equal weight of liquid. If gas is compressed to  $1/1600^{\text{th}}$  of its volume above critical temperature (which is different for each gas) no phase change occurs and the resulting substance is a gas that is just as dense as a liquid.


### Gas Laws

- **Boyle's law** : It states that at a constant temperature ( $T$ ),  $P \propto (1/V)$  or  $PV = \text{constant} \Rightarrow P_1 V_1 = P_2 V_2$
- **Charles' law** : It states that at constant pressure ( $P$ ),  $V \propto T$  or  $(V/T) = \text{constant} \Rightarrow V_1/T_1 = V_2/T_2$
- **Avogadro law** : It states that at constant temperature ( $T$ ) and pressure ( $P$ ),  $V \propto n$  or  $(V/n) = \text{constant}$
- **Gay Lussac's law** : It states that at constant volume ( $V$ ),  $P \propto T$  or  $(P/T) = \text{constant}$ .

#### Boyle's Law



#### Charles' Law



#### Ideal Gas Equation

$$PV = nRT \quad [R = \text{gas constant}]$$

$$0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1} \quad \boxed{R \text{ (gas constant)}} \quad 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1} \\ 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \quad 1.99 \text{ cal K}^{-1} \text{ mol}^{-1}$$


#### Deviation from Ideal Gas Behaviour (Real Gas)

- Real gas follows gas laws only at high temperature and low pressure.
- It follows van der Waals' equation;  $\left(P + \frac{an^2}{V^2}\right)(V - nb) = nRT$

#### Compressibility Factor

$Z = 1$  refers to ideal behaviour  
 $Z > 1$  shows positive deviation  
 $Z < 1$  shows negative deviation

where,  $Z$  is compressibility factor which measures the deviation from ideal behaviour.



- **Relation between critical constants and van der Waals' constants**:  $V_c = 3b$ ,  $P_c = \frac{a}{27b^2}$ ,  $T_c = \frac{8a}{27Rb}$
  - As the value of  $a$  increases, the critical temperature ( $T_c$ ) also increases. Higher the critical temperature of the gas, more easily it liquefies.
- The ease of liquefaction order:  $\text{Cl}_2 > \text{CO}_2 > \text{CH}_4 > \text{H}_2 > \text{He}$

### Graham's Law of Diffusion

$$\frac{r_1}{r_2} = \frac{v_1/t_1}{v_2/t_2} = \sqrt{\frac{d_2}{d_1}} = \sqrt{\frac{M_2}{M_1}}$$

### Dalton's Law of Partial Pressure

It states that "at a given temperature, the total pressure ( $P_{\text{total}}$ ) exerted by two or more non-reacting gases occupying a definite volume is equal to the sum of partial pressures of the component gases".

$$P_{\text{total}} = p_1 + p_2 + p_3 + \dots \quad (\text{where } p_1, p_2, p_3 \text{ are partial pressures of component gases})$$


### Kinetic Molecular Theory of Gases

- All gases are made up of a very large number of minute particles like atoms or molecules.
- The molecules are in continuous random motion in all directions.
- Molecular collisions are perfectly elastic.
- The intermolecular forces of attraction are negligible.
- The average kinetic energy of a gas molecule is directly proportional to the absolute temperature.
- Kinetic gas equation:  $PV = \frac{1}{3} mnu^2$

### Molecular Speed

- Root mean square velocity :  $u_{rms} = \sqrt{\frac{3RT}{M}}$
  - Average velocity :  $u_{av} = \sqrt{\frac{8RT}{\pi M}}$
  - Most probable speed :  $u_{mp} = \sqrt{\frac{2RT}{M}}$
- $$u_{mp} : u_{av} : u_{rms} = 1 : 1.128 : 1.224$$

### Maxwell-Boltzmann Distribution Curve



Signifies that speed is never zero but it tends towards zero.

# Diazonium Salt

**General formula :**  $\text{ArN}_2^+\text{X}^-$   
 $\text{X}^-$  may be  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{HSO}_4^-$ ,  $\text{BF}_4^-$ , etc.


Azo dyes can be prepared using diazonium salts, these contain a highly delocalised system of electrons which delocalise in both benzene rings and the two nitrogen atoms bridging the rings. This extended delocalisation give colours to these azo compounds. Modifying the group present in molecule can give rise to different colours.

## CONCEPT MAP

CLASS XII

### Preparation

The reaction of converting aromatic primary amine to diazonium salt is called diazotisation.



### Physical Properties


- Colourless crystalline solids.
- Readily soluble in water, less soluble in alcohol.
- These are unstable and explode in dry state.
- Their aqueous solutions are neutral to litmus.

### Applications

- To produce water-fast dyed fabrics by immersing the fabric in an aqueous solution of the diazonium compound, followed by immersion in a solution of the coupler.
- Synthesis of organic compounds.
- Diazonium salts are light sensitive and break down near UV or violet light. This property has led to their use in document reproduction.







### Chemical Properties

#### Reactions Involving Displacement of Diazo Group



#### Reactions Involving Retention of Diazo group

- The azo products obtained have an extended conjugate system having both aromatic rings joined through  $-\text{N}=\text{N}-$  bond.
- These compounds are often coloured and are used as a dye.







## PHYSICAL PROPERTIES


- Lower alcohols are liquid at room temperature while higher ones are solid.
- Solubility in water decreases with increase in molecular mass due to decrease in extent of intermolecular hydrogen bonding.
- Alcohols have high boiling points. Order of boiling point : 1° alcohols > 2° alcohols > 3° alcohols

## CHEMICAL PROPERTIES


- The hydroxyl group present in alcohols is a very reactive group. The reactions of the hydroxyl group consists of either cleavage of C—O bond or the cleavage of O—H bond as both are highly polarised with negative end of the dipoles at oxygen.
- In O—H cleavage, the order of reactivity : Primary > Secondary > Tertiary
- In C—O cleavage, the order of reactivity : Tertiary > Secondary > Primary

	$\xrightarrow{\text{Na}} \text{RONa} + \text{H}_2$
<b>A</b>	$\xrightarrow[\text{Conc. H}_2\text{SO}_4]{\text{RCOOH}} \text{RCOOR} + \text{H}_2\text{O}$
	$\xrightarrow{\text{RCOCl}} \text{RCOOR} + \text{HCl}$
<b>L</b>	$\xrightarrow[\text{Base}]{(\text{RCO})_2\text{O}} \text{RCOOR} + \text{RCOOH}$
<b>C</b>	$\xrightarrow{\text{R'MgX}} \text{R'H} + \text{Mg(OR)X}$
	$\xrightarrow{\text{HX}} \text{RX} + \text{H}_2\text{O}$
<b>O</b>	$\xrightarrow{\text{PX}_3} \text{RX} + \text{H}_3\text{PO}_3$
	$\xrightarrow{\text{PCl}_5} \text{RCl} + \text{POCl}_3 + \text{HCl}$
<b>H</b>	$\xrightarrow{\text{SOCl}_2} \text{RCl} + \text{SO}_2\uparrow + \text{HCl}\uparrow$
	$\xrightarrow[443\text{ K}]{\text{Conc. H}_2\text{SO}_4} \text{RCH=CH}_2 + \text{H}_2\text{O}$
<b>O</b>	$\xrightarrow[413\text{ K}]{\text{Conc. H}_2\text{SO}_4} \text{ROR} + \text{H}_2\text{O}$
<b>L</b>	$\xrightarrow[\text{or KMnO}_4/\text{OH}^-]{\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+} \text{R-COOH}$
	$\xrightarrow[\text{-H}_2]{\text{Cu/573 K}} \text{RCHO or RCOR or alkene}$ (from 1° alcohol) (from 2° alcohol) (from 3° alcohol)
<b>S</b>	$\xrightarrow[\text{or CrO}_3\cdot\text{C}_5\text{H}_5\text{N}\cdot\text{HCl}]{\text{PCC}} \text{RCHO}$

## Dehydration of alcohols :



### Mechanism :




- The order of stability of carbocations : 3° > 2° > 1°
- It always occurs in accordance with the Saytzeff rule *i.e.*, the more substituted alkene is the major product.

## Distinction between 1°, 2° and 3° Alcohols

- Lucas test :** Alcohols +  $\text{ZnCl}_2 + \text{HCl}$ 
  - 1° Alcohol :  $\text{RCH}_2\text{OH} + \text{ZnCl}_2 + \text{HCl} \longrightarrow$   
No reaction at room temperature
  - 2° Alcohol :  $\text{R}_2\text{CHOH} + \text{ZnCl}_2 \longrightarrow \text{R}_2\text{CHCl}$ ;  
White turbidity appeared after 5-10 min.
  - 3° Alcohol :  $\text{R}_3\text{COH} + \text{ZnCl}_2 + \text{HCl} \longrightarrow \text{R}_3\text{CCl}$ ;  
White turbidity appeared immediately.


### Victor Meyer test :

#### 1° Alcohol :




Nitrolic acid on treatment with alkali gives blood red colouration.

#### 2° Alcohol :



Pseudonitrol on treatment with alkali gives blue colouration.

➤ **3° Alcohol :**




## PHENOLS


- Phenols are derivatives of benzene in which a ring hydrogen is replaced by —OH group.

## PREPARATION



- Hydrolysis of chlorobenzene (Dow's process) :**



- From benzene sulphonic acid (alkali fusion of sodium benzene sulfonate) :**




- From cumene hydroperoxide :**



## PHYSICAL PROPERTIES


- Phenol is a colourless, crystalline, deliquescent solid, attains pink colour on exposure to air and light.
- Its melting point is 315 K and boiling point 455 K. The boiling point of phenol is much higher than the corresponding aromatic hydrocarbons and the haloarenes.
- It is soluble in water due to inter-molecular H-bonding among themselves and with water.
- It is poisonous in nature but acts as antiseptic and disinfectant.

## CHEMICAL PROPERTIES




• **Test for phenols :**

- **Ferric chloride test :** Phenol gives violet colour with neutral  $\text{FeCl}_3$  solution.



- **Bromine water test :** Phenol gives white ppt. with  $\text{Br}_2$ -water due to the formation of 2, 4, 6-tribromophenol.



- **Acidity of phenol :** Phenols are much more acidic than alcohols but less than carboxylic acids or even carbonic acid. This is indicated by the values of ionisation constants ( $K_a$ ) or  $\text{p}K_a$ . Smaller the  $\text{p}K_a$  value, stronger the acid. The relative acidity follows the following order:


	$K_a$ (approx.)	$\text{p}K_a$ value
Acidic strength decreases	$(10^{-5})$ $\text{RCOOH}$ Carboxylic acid	5
	$(10^{-7})$ $\text{H}_2\text{CO}_3$ Carbonic acid	7
	$(10^{-10})$ $\text{C}_6\text{H}_5\text{OH}$ Phenol	8-10
	$(10^{-14})$ $\text{HOH}$ Water	14
	$(10^{-18})$ $\text{ROH}$ Alcohols	16-18

## ETHERS


- **General formula :**  $\text{C}_n\text{H}_{2n+2}\text{O}$  ( $\text{R}-\text{O}-\text{R}'$ )
- **Symmetrical ether :** When  $\text{R}$  and  $\text{R}'$  are same groups. e.g.,  $\text{CH}_3-\text{O}-\text{CH}_3$ ,  $\text{C}_2\text{H}_5-\text{O}-\text{C}_2\text{H}_5$
- **Unsymmetrical ether :** When  $\text{R}$  and  $\text{R}'$  are different groups. e.g.,  $\text{CH}_3-\text{O}-\text{C}_6\text{H}_5$ ,  $\text{CH}_3-\text{O}-\text{C}_2\text{H}_5$

## PREPARATION


### From alcohols by dehydration :




### From alkyl halides :




### From Grignard reagent :



### From diazomethane :



### From alkenes :




## PHYSICAL PROPERTIES



- Ethers have a bent structure and are dipolar in nature.
- Boiling points of ethers show a gradual increase with increase in molecular mass.
- They are lighter than water. Lower ethers are highly volatile and very inflammable.
- They are sparingly soluble in water but readily soluble in organic solvents.
- Dimethyl ether and ethyl methyl ether are gases. All ethers are colourless liquids with characteristic ether smell.

## CHEMICAL PROPERTIES





- **Reactions of lone pair of oxygen :**




• **Cleavage of R—O—R bond :**



• **Different possibilities of reaction of ether with HI :**



• **Electrophilic substitution reactions of aromatic ethers :**



**USES**

- Diethyl ether is used as solvent for oils, fats, waxes, plastics and lacquers.
- Provide inert medium for reactions.
- Used in surgery as an anaesthetic.
- Used in perfumery and in the manufacture of smokeless powder.

**INSIGHTS**

**A bioinspired *ortho*-sulfiliminyl phenol synthesis!**


A variety of synthetic methods have been developed to construct the *ortho*-functionalized phenols which are highly useful in chemical industry, functional materials and medicines. These methods mainly include three kind of strategies (a) rearrangement of aromatic O—X bonds (b) directing group-assisted *ortho* C—H hydroxylation of arenes and (c) *ortho* C—H functionalization of phenols.



# SPEED PRACTICE


1. Which one of the following properties is exhibited by phenol?
- It is soluble in aq. NaOH and evolves  $\text{CO}_2$  with aq.  $\text{NaHCO}_3$ .
  - It is soluble in aq. NaOH and does not evolve  $\text{CO}_2$  with aq.  $\text{NaHCO}_3$ .
  - It is not soluble in aq. NaOH but evolves  $\text{CO}_2$  with aq.  $\text{NaHCO}_3$ .
  - It is insoluble in aq. NaOH and does not evolve  $\text{CO}_2$  with aq.  $\text{NaHCO}_3$ .

2. The major product of the following reaction is



- $$\begin{array}{c} \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2 \diagdown \\ \text{CH}_3 \diagup \text{C}=\text{CH}_2 \end{array}$$
- $$\begin{array}{c} \text{C}_6\text{H}_5 \diagdown \quad \text{H} \diagup \\ \text{H} \diagup \text{C}=\text{C} \diagdown \\ \quad \quad \quad \text{CH}(\text{CH}_3)_2 \end{array}$$
- $$\begin{array}{c} \text{C}_6\text{H}_5\text{CH}_2 \diagdown \quad \text{CH}_3 \diagup \\ \text{H} \diagup \text{C}=\text{C} \diagdown \\ \quad \quad \quad \text{CH}_3 \end{array}$$
- $$\begin{array}{c} \text{C}_6\text{H}_5 \diagdown \quad \text{CH}(\text{CH}_3)_2 \diagup \\ \text{H} \diagup \text{C}=\text{C} \diagdown \\ \quad \quad \quad \text{H} \end{array}$$

3. The reaction





can be classified as

- dehydration reaction
- Williamson alcohol synthesis reaction
- Williamson ether synthesis reaction
- alcohol formation reaction.

(NEET Phase-I 2016)

4. Identify the set of reagents/reaction conditions, (x and y) in the following set of transformation.



- $x$  = dilute aqueous NaOH,  $20^\circ\text{C}$ ;  $y$  = HBr/acetic acid,  $20^\circ\text{C}$

(b)  $x$  = conc. alcoholic NaOH,  $80^\circ\text{C}$ ;  $y$  = HBr/acetic acid,  $20^\circ\text{C}$

(c)  $x$  = dilute aqueous NaOH,  $20^\circ\text{C}$ ;  $y$  =  $\text{Br}_2/\text{CHCl}_3$ ,  $0^\circ\text{C}$

(d)  $x$  = conc. alcoholic NaOH,  $80^\circ\text{C}$ ;  $y$  =  $\text{Br}_2/\text{CHCl}_3$ ,  $0^\circ\text{C}$

5.  $\text{CH}_3\text{Br} + \text{Nu}^- \longrightarrow \text{CH}_3 - \text{Nu} + \text{Br}^-$

The decreasing order of the rate of the above reaction with the given nucleophiles is

$[\text{Nu}^- = (\text{A}) \text{PhO}^-, (\text{B}) \text{AcO}^-, (\text{C}) \text{HO}^-, (\text{D}) \text{CH}_3\text{O}^-]$

- $\text{D} > \text{C} > \text{A} > \text{B}$
- $\text{D} > \text{C} > \text{B} > \text{A}$
- $\text{A} > \text{B} > \text{C} > \text{D}$
- $\text{B} > \text{D} > \text{C} > \text{A}$


6. Etherates are

- ethers
- solution in ether
- complexes of ethers with Lewis acids
- complexes of ethers with Lewis bases.

7. An organic compound (X) with molecular formula  $\text{C}_7\text{H}_8\text{O}$  is insoluble in aqueous  $\text{NaHCO}_3$  but dissolves in NaOH. When treated with bromine water (X) rapidly gives (Y),  $\text{C}_7\text{H}_5\text{OBr}_3$ . The compounds (X) and (Y) respectively are



- benzyl alcohol and 2, 4, 6-tribromo-3-methoxy benzene
- benzyl alcohol and 2, 4, 6-tribromo-3-methyl phenol
- o*-cresol and 3, 4, 5-tribromo-2-methyl phenol
- m*-cresol and 2, 4, 6-tribromo-3-methyl phenol.

8.  $\text{A} \xrightarrow[\text{One. eq. of CH}_3\text{I}]{\text{OH}^-} \text{C}_9\text{H}_{10}\text{O}_3 \xrightarrow[\text{(HCO}_3^- \text{ soluble)}]{\text{KMnO}_4} \text{C}_9\text{H}_{10}\text{O}_4$




Starting substrate 'A' (gives intense colour with  $\text{FeCl}_3$  and positive Tollens' test) is


- 
-



The compounds A, B and C respectively are




10.  $S_N1$  reaction undergoes through a carbocation intermediate as follows :




Choose the correct statements for  $S_N1$  reaction.

- (I) The decreasing order of rate of  $S_N1$  reaction is  $t\text{-BuX} > \text{iso-PrX} > \text{EtX} > \text{MeX}$
- (II) The decreasing order of ionization energy is  $\text{MeX} > \text{EtX} > \text{iso-PrX} > t\text{-BuX}$ .
- (III) The decreasing order of energy of activation is  $t\text{-BuX} > \text{iso-PrX} > \text{EtX} > \text{MeX}$ .
- (a) Only I and II      (b) Only I and III  
(c) Only II and III    (d) All of these.

11. Identify A and predict the type of reaction.



(NEET 2017)




Here, B, C and D respectively are

- (a) aspirin, salol, oil of winter green  
(b) salol, aspirin, oil of winter green  
(c) oil of winter green, aspirin, salol  
(d) oil of winter green, salol, aspirin.

13. Chloropicrin is obtained by the reaction of

- (a) chlorine on picric acid  
(b) nitric acid on chloroform  
(c) steam on carbon tetrachloride  
(d) nitric acid on chlorobenzene.

14. For the following compounds, the correct statement(s) with respect to nucleophilic substitution reactions is (are)





- (a) I and II follow  $S_N2$  mechanism  
(b) compound IV undergoes inversion of configuration

- (c) the order of reactivity for I, III, and IV is :  
IV > I > III  
(d) I and III follow  $S_N1$  mechanism.


[JEE Advanced 2017]

15. Consider thiolate ion ( $RS^-$ ) and alkoxide ion ( $RO^-$ ). Which of the following statements is correct?  
(a)  $RS^-$  is more basic and less nucleophilic than  $RO^-$ .  
(b)  $RS^-$  is less basic and less nucleophilic than  $RO^-$ .  
(c)  $RS^-$  is less basic but more nucleophilic than  $RO^-$ .  
(d)  $RS^-$  is more basic and more nucleophilic than  $RO^-$ .
16. Diethyl ether when refluxed with excess of HI gives two molecules of (i). Ethers can be most commonly prepared by reaction of (ii) and (iii). The method is called (iv).  
Here, (i), (ii), (iii) and (iv) respectively are  
(a) ethyl iodide, sodium alkoxide, alkyl halide, Williamson synthesis  
(b) ethanol, alcohol, alkyl halide, substitution  
(c) methyl iodide, Grignard reagent, alkyl halide, Williamson synthesis  
(d) ethyl iodide, phenol, alkyl iodide, esterification.
17. Phenyl methyl ether (anisole) reacts with HI to give phenol and methyl iodide but not iodobenzene and methyl alcohol because  
(a)  $I^-$  attacks the methyl group as C—O bond between phenyl and oxygen is formed by  $sp^2$  hybridised carbon, thus difficult to break  
(b)  $I^-$  ion is not reactive towards benzene  
(c) phenol is formed as a result of hydrolysis of iodobenzene  
(d) methyl alcohol formed during reaction, reacts with  $I^-$  to form methyl iodide.
18. Consider the following reaction,  
 $CH_3CH_2CH_2Br + NaCN \rightarrow CH_3CH_2CH_2CN + NaBr$   
This reaction will be the fastest in  
(a) ethanol (b) methanol  
(c) *N,N*-dimethylformamide (DMF)  
(d) water. [NEET Phase-II 2016]


19. In the dehydrohalogenation of 2-bromobutane, which conformation leads to the formation of *cis*-2-butene?



20. In the following series of chemical reactions, identify Z :



- (a)  $CH_3-\underset{\text{Br}}{\underset{|}{CH}}-\underset{\text{OH}}{\underset{|}{CH_2}}$  (b)  $CH_3-\underset{\text{OH}}{\underset{|}{CH}}-\underset{\text{OH}}{\underset{|}{CH_2}}$   
(c)  $CH_3-\underset{\text{Br}}{\underset{|}{C}}=CH_2$  (d)  $CH_3C \equiv CH$
21. Glycerol forms salt with sodium metal at 110 °C. The sodium salt is  
(a) monosalt (b) disalt  
(c) trisalt (d) mixture of monosalt and disalt.
22. In the following reaction sequence :




The compound I is


- (a)  $\begin{array}{c} CH_2-CH-CH_3 \\ | \quad | \\ Cl \quad Cl \end{array}$   
(b)  $\begin{array}{c} Cl \\ | \\ CH_3-C-CH_3 \\ | \\ Cl \end{array}$   
(c)  $\begin{array}{c} Cl-CH-CH_2-CH_3 \\ | \\ Cl \end{array}$   
(d)  $\begin{array}{c} CH_2-CH_2-CH_2 \\ | \quad \quad | \\ Cl \quad \quad Cl \end{array}$

[JEE Main Online 2017]

23. Dehydrohalogenation by strong base is slowest in




24. In the following compounds, nucleophile and the leaving groups are present in the same molecule :




These dual nature species can undergo intramolecular and intermolecular nucleophilic substitution reactions. Intramolecular substitution reaction is possible only in

- (a) I, II (b) II, III (c) III, IV (d) IV
25. 0.037 g of an alcohol,  $R-OH$  was added to  $CH_3MgBr$  and the gas evolved measured 11.2 mL at STP. The molecular mass of  $R-OH$  will be
- (a) 47 (b) 79 (c) 74 (d) 77



[JEE Main 2016]

27. Which of the following reagents cannot be used to oxidise primary alcohols to aldehydes?
- (a)  $CrO_3$  in anhydrous medium  
 (b)  $KMnO_4$  in acidic medium  
 (c) Pyridinium chlorochromate  
 (d) Heat in the presence of Cu at 573 K
28. Chloromethane on treatment with excess of ammonia yields mainly
- (a)  $N,N$ -dimethylmethanamine  
 (b)  $N$ -methylmethanamine  
 (c) methanamine  
 (d) mixture containing all these in equal proportion.




- (a) (1R, 3S)-cis-3-methylcyclohexanol  
 (b) (1S, 3S)-cis-3-methylcyclohexanol  
 (c) (1S, 3S)-trans-3-methylcyclohexanol  
 (d) (1R, 3R)-trans-3-methylcyclohexanol.

30. Bottles containing  $C_6H_5I$  and  $C_6H_5CH_2I$  lost their original labels. They were labelled A and B for testing. A and B were separately taken in test tubes and boiled with NaOH solution. The end solution in each tube was made acidic with dilute  $HNO_3$  and some  $AgNO_3$  solution added. Solution B gave yellow precipitate. Which one of the following statements is true for the experiment?
- (a) Addition of  $HNO_3$  was unnecessary.  
 (b) A was  $C_6H_5I$ .  
 (c) A was  $C_6H_5CH_2I$ .  
 (d) B was  $C_6H_5I$ .


## SOLUTIONS

1. (b) : Phenol is soluble in aq. NaOH forming sodium phenoxide.  
 $C_6H_5OH + NaOH \longrightarrow C_6H_5ONa + H_2O$   
 But phenol does not evolve  $CO_2$  with aq.  $NaHCO_3$  and aq.  $Na_2CO_3$  because phenol is weaker acid than carbonic acid.




Trans is major product as it is thermodynamically more stable.

3. (c) 4. (b)
5. (a) :  $CH_3O^- > HO^- > \text{Phenoxide}^- > CH_3COO^-$   
 (D) (C) (A) (B)







9. (b)

10. (a): For  $S_N1$  reaction, rate depends upon the stability of carbocation formed. Thus, rate of  $S_N1$  reaction is  $t\text{-BuX} > \text{iso-PrX} > \text{EtX} > \text{MeX}$

Higher rate of reaction is possible only if activation energy is low.

So, order of activation energy is




Order of ionization energy is



11. (d): *m*-Bromoanisole gives only the respective *meta*-substituted product through benzyne intermediate. This is a substitution reaction which goes by an elimination-addition pathway.

12. (a)



14. (a, b, d):

(a) Compounds I and II are  $1^\circ$  alkyl halides, thus they undergo  $S_N2$  mechanism.

(b) Compound IV undergoes inversion of configuration due to intimate-ion pair formation, inversion predominates over retention.

(c) Stability of carbocations follows the order:  
 $2^\circ$  benzylic (IV)  $>$   $3^\circ$  alkyl (III)  $>$   $1^\circ$  benzylic (I)

(d) I is a benzylic halide, thus, it also undergoes  $S_N1$  reaction easily as benzylic carbocation is resonance stabilised and III also follows  $S_N1$  mechanism as it is a  $3^\circ$  alkyl halide.


15. (c)

16. (a)


17. (a)

18. (c)

19. (a)




21. (d)



23. (c)

24. (c)




11.2 mL  $\text{CH}_4$  at STP is formed from 0.037 g ROH.

$\therefore$  22400 mL  $\text{CH}_4$  at STP will be formed from

$$\frac{0.037}{11.2} \times 22400 \text{ g ROH} = 74 \text{ g ROH}$$


$\therefore$  Molecular mass of ROH = 74



27. (b)

28. (c)


29. (c):



30. (b)



# EXAMINER'S MIND CLASS XII



The questions given in this column have been prepared strictly on the basis of NCERT Chemistry for Class XII. This year JEE (Main & Advanced)/NEET/AIIMS have drawn their papers heavily from NCERT books.

<b>Section - I</b>	Q. 1 to 10 Only One Option Correct Type MCQs.
<b>Section - II</b>	Q. 11 to 13 More than One Options Correct Type MCQs.
<b>Section - III</b>	Q. 14 to 17 Paragraph Type MCQs having Only One Option Correct.
<b>Section - IV</b>	Q. 18 & 19 Matching List Type MCQs having Only One Option Correct.
<b>Section - V</b>	Q. 20 to 22 Assertion Reason Type MCQs having Only One Option Correct. Mark the correct choice as : (a) If both assertion and reason are true and reason is the correct explanation of assertion. (b) If both assertion and reason are true but reason is not the correct explanation of assertion. (c) If assertion is true but reason is false. (d) If both assertion and reason are false.
<b>Section - VI</b>	Q. 23 to 25 Integer Value Correct Type Questions having Single Digit Integer Answer, ranging from 0 to 9 (both inclusive).


## ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

### SECTION - I

#### Only One Option Correct Type

- Which of the following is paraldehyde?  
(a)  $(\text{HCHO})_4$  (b)  $(\text{CH}_3\text{CHO})_3$   
(c)  $(\text{HCHO})_3$  (d)  $(\text{CH}_3\text{CHO})_4$
- The carboxyl functional group is present in  
(a) picric acid (b) barbituric acid  
(c) ascorbic acid (d) aspirin.
- Identify the product Y in the following reaction sequence :  

$$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{COO} \\ | \\ \text{CH}_2-\text{CH}_2-\text{COO} \end{array} \text{Ca} \xrightarrow{\text{heat}} \text{X} \xrightarrow[\text{conc. HCl}]{\text{Zn-Hg}} \text{Y}$$
- Identify the starting reagents (X and Y) needed to make the following compound by crossed aldol condensation.




- Acetophenone and butanal  
(b) Benzaldehyde and 2-pentanone  
(c) Acetophenone and 2-butanone  
(d) Benzaldehyde and 3-pentanone
- In an ester molecule there are three C — O bonds,  


$$\text{R}-\overset{\gamma}{\underset{\text{O}}{\underset{||}{\text{C}}}}-\overset{\beta}{\text{O}}-\overset{\alpha}{\text{R}'}$$
What do you expect regarding their relative bond lengths?  
(a)  $\alpha < \beta < \gamma$  (b)  $\beta = \gamma > \alpha$   
(c)  $\beta > \gamma > \alpha$  (d)  $\gamma < \beta < \alpha$
- In order to prepare acetone from acetyl chloride in one step, which of the following reagents will be the best?  
(a) Reduction with  $\text{H}_2/\text{Pd}-\text{BaSO}_4$   
(b) Reduction with HI  
(c) Grignard reagent  
(d) Dimethyl cadmium
- $2\text{CH} \equiv \text{CH} \xrightarrow[\text{H}_2\text{SO}_4]{\text{HgSO}_4} (\text{X}) \xrightarrow{\text{LiAlH}_4} (\text{Y}) \xrightarrow{\text{P/Br}_2} (\text{Z})$   
In the given sequence of reactions, (Z) is  
(a) ethylene bromide (b) ethanol  
(c) ethyl bromide (d) ethylidene bromide.

8. IUPAC name of the following compound is

- (a) heptane 2, 6-dione  
(b) octan-6-one  
(c) octane 2, 7-dione  
(d) hexane-2, 5-dione.



9. In the following reaction sequence, the correct structures of A, B and C are



- (a)  $\text{A} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$      $\text{B} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{ONa}$      $\text{C} = \text{CHI}_3$   
 (b)  $\text{A} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$      $\text{B} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{ONa}$      $\text{C} = \text{CHI}_3$   
 (c)  $\text{A} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$      $\text{B} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{ONa}$      $\text{C} = \overset{*}{\text{C}}\text{HI}_3$   
 (d)  $\text{A} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{CH}_3$      $\text{B} = \text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\overset{*}{\text{C}}-\text{ONa}$      $\text{C} = \overset{*}{\text{C}}\text{H}_3\text{I}$

10. The correct sequence of steps involved in the mechanism of Cannizzaro reaction is

- (a) nucleophilic attack, transfer of  $\text{H}^-$  and transfer of  $\text{H}^+$   
 (b) transfer of  $\text{H}^-$ , transfer of  $\text{H}^+$  and nucleophilic attack  
 (c) transfer of  $\text{H}^+$ , nucleophilic attack and transfer of  $\text{H}^-$   
 (d) electrophilic attack by  $\text{OH}^-$ , transfer of  $\text{H}^+$  and transfer of  $\text{H}^-$ .

## SECTION - II

More than One Options Correct Type

11. Reaction of  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$  with a mixture of  $\text{Br}_2$  and  $\text{KOH}$  gives  $\text{R}-\text{NH}_2$  as the main product. The intermediates involved in this reaction are

- (a)  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NHBr}$     (b)  $\text{R}-\text{NHBr}$   
 (c)  $\text{R}-\text{N}=\text{C}=\text{O}$     (d)  $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{N}(\text{Br})_2$

12. Which of the following reactions is used for detecting the presence of carbonyl group?

- (a) Reaction with hydroxylamine  
 (b) Reaction with hydrazine  
 (c) Reaction with phenyl hydrazine  
 (d) Reaction with semicarbazide

13. Aqueous solution of sodium succinate on electrolysis gives

- (a)  $\text{CH}_3-\text{CH}_3$     (b)  $\text{CH}_2=\text{CH}_2$   
 (c)  $\text{CH}\equiv\text{CH}$     (d)  $\text{CO}_2$

## SECTION - III

Paragraph Type

### Paragraph for Questions 14 and 15

The acidic strength of saturated aliphatic carboxylic acids depends mainly upon the inductive effect of the substituent and its position with respect to the  $-\text{COOH}$  group. Electron donating substituents tend to decrease, whereas electron withdrawing substituents tend to increase the acidic strength. The effect of electron donating substituents and electron withdrawing substituents is more pronounced at *p*-position than that at *m*-position. Due to *ortho* effect, *o*-substituted benzoic acids are usually stronger than benzoic acid regardless of the nature of substituent whether electron donating or electron withdrawing.

14. Which of the following is expected to be highly ionized in water?


- (a)  $\text{ClCH}_2\text{CH}_2\text{CH}_2\text{COOH}$   
 (b)  $\text{CH}_3\text{CHClCH}_2\text{COOH}$   
 (c)  $\text{CH}_3\text{CH}_2\text{CCl}_2\text{COOH}$   
 (d)  $\text{CH}_3\text{CH}_2\text{CHClCOOH}$

15. Which of the following acids has the smallest dissociation constant?

- (a)  $\text{CH}_3\text{CHFCH}_2\text{COOH}$     (b)  $\text{FCH}_2\text{CH}_2\text{COOH}$   
 (c)  $\text{BrCH}_2\text{CH}_2\text{COOH}$     (d)  $\text{CH}_3\text{CHBrCH}_2\text{COOH}$

### Paragraph for Questions 16 and 17

In the following reaction sequence, the compound J is an intermediate.




J( $\text{C}_9\text{H}_8\text{O}_2$ ) gives effervescence on treatment with  $\text{NaHCO}_3$  and positive Baeyer's test.

16. The compound K is

- (a)    (b)   
 (c)    (d)

17. The compound I is



#### SECTION - IV


##### Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List I		List II	
(P) Schiff's reagent	(i) Elimination reaction		
(Q) Alcoholic KOH	(ii) Colourless solution due to $\text{SO}_2$		
(R) Ammoniacal $\text{AgNO}_3$	(iii) Aldehydes		
(S) Benedict solution	(iv) Chloroform		

P	Q	R	S
(a) (ii) (iii)	(i)	(iii) (iv)	(iii)
(b) (ii) (iii)	(ii)	(ii) (i)	(ii) (iv)
(c) (ii)	(i)	(iii)	(iv)
(d) (i)	(ii)	(i) (iii)	(iii) (iv)

19. Match the List I with List II and select the correct answer using the codes given below the lists :




P	Q	R	S
(a) (i)	(ii)	(iii)	(iv)
(b) (iii)	(iv)	(i)	(ii)
(c) (iv)	(i)	(ii)	(iii)
(d) (ii)	(iii)	(iv)	(i)

#### SECTION - V

##### Assertion Reason Type

20. **Assertion :** The following reaction occurs



**Reason :** The semicarbazone of (A) does not show geometrical isomerism.

21. **Assertion :** Esters which contain  $\alpha$ -hydrogens undergo Claisen condensation reaction.


**Reasons :** Reduction of esters by  $\text{LiAlH}_4$  gives acids.

22. **Assertion :**  $\text{HCHO}$  reacts with  $\text{NH}_3$  to give urotropine.

**Reason :** Urotropine is a medicine for troubles in urinary track.

#### SECTION - VI


##### Integer Value Correct Type



How many hydrogen atoms are replaced by D?

24. If formic acid, acetic acid, propanoic acid and benzoic acid are mixed with phosphorus and bromine then how many products will be formed?

25. The number of compounds among the following which give iodoform test is



#### Your favourite MTG Books/Magazines available in PONDICHERRY at

- Siruvur Pakkam - Pondicherry Ph: 2338686; Mob: 9894034367, 9087734367
- Sri Sivagami News Agency - Pondicherry Mob: 9443074953

Visit "MTG IN YOUR CITY" on [www.mtg.in](http://www.mtg.in) to locate nearest book seller OR write to [info@mtg.in](mailto:info@mtg.in) OR call 0124-6601200 for further assistance.



## COORDINATION COMPOUNDS

### SECTION - I

#### Only One Option Correct Type

- Crystal field stabilisation energy for high spin  $d^4$  octahedral complex is  
(a)  $-0.6\Delta_o$  (b)  $-1.8\Delta_o$   
(c)  $-1.6\Delta_o + P$  (d)  $-1.2\Delta_o$
- A solution containing 2.675 g of  $\text{CoCl}_3 \cdot 6\text{NH}_3$  (molar mass =  $267.5 \text{ g mol}^{-1}$ ) is passed through a cation exchanger. The chloride ions obtained in the solution were treated with excess of  $\text{AgNO}_3$  to give 4.32 g of  $\text{AgCl}$  (molar mass =  $143.5 \text{ g mol}^{-1}$ ). The formula of the complex is (At. mass of Ag = 108 u)  
(a)  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$   
(b)  $[\text{CoCl}_2(\text{NH}_3)_4]\text{Cl} \cdot 2\text{NH}_3$   
(c)  $[\text{CoCl}_3(\text{NH}_3)_3] \cdot 3\text{NH}_3$   
(d)  $[\text{CoCl}(\text{NH}_3)_5]\text{Cl}_2 \cdot \text{NH}_3$
- According to IUPAC nomenclature, sodium nitroprusside is named as  
(a) sodium nitroferrocyanide  
(b) sodium nitroferrocyanide  
(c) sodium pentacyanonitrosylferrate (II)  
(d) sodium pentacyanonitrosylferrate (III).
- Ferrocene is an example of  
(a) sandwiched complex (b) pi-bonded complex  
(c) a complex in which all the five carbon atoms of cyclopentadiene anions are bonded to the metal  
(d) all of the above.
- Which kind of isomerism is exhibited by octahedral complex,  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$ ?  
(a) Geometrical and ionisation  
(b) Geometrical and optical  
(c) Optical and ionisation  
(d) Geometrical only
- When EDTA solution is added to  $\text{Mg}^{2+}$  ion solution, then which of the following statements is not true?  
(a) Four coordinate sites of  $\text{Mg}^{2+}$  are occupied by EDTA and remaining two sites are occupied by water molecules.  
(b) All six coordinate sites of  $\text{Mg}^{2+}$  are occupied.  
(c) pH of the solution is decreased.  
(d) Colourless  $[\text{Mg} - \text{EDTA}]^{2-}$  chelate is formed.
- The oxidation number,  $d$ -orbital occupation and coordination number of Cr in the complex  $\text{cis-}[\text{Cr}(\text{en})_2\text{Cl}_2]\text{Cl}$  are respectively  
(a) +3,  $3d$  and 4 (b) +3,  $4d$  and 6  
(c) +3,  $3d$  and 6 (d) +2,  $3d$  and 6

8. If  $P > \Delta_o$  then  $d^4$  is represented as

- (a)  $t_{2g}^{211} e_g^0$  (b)  $t_{2g}^{111} e_g^1$   
(c)  $t_{2g}^0 e_g^{22}$  (d)  $t_{2g}^1 e_g^{21}$

9. What are the correct oxidation state, coordination number, configuration, magnetic character and magnetic moment of  $\text{K}_4[\text{Mn}(\text{CN})_6]$ ?

	O.S.	C.N.	Confi- guration	Magnetic Character	Magnetic Moment
(a)	+6	6	$t_{2g}^5 e_g^0$	Diamagnetic	0
(b)	+2	6	$t_{2g}^4 e_g^1$	Diamagnetic	0
(c)	+2	6	$t_{2g}^5 e_g^0$	Paramagnetic	1.732
(d)	+4	6	$t_{2g}^3 e_g^2$	Diamagnetic	0


10. Which of the following isomers will give white precipitate with  $\text{BaCl}_2$  solution?

- (a)  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$  (b)  $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$   
(c)  $[\text{Co}(\text{NH}_3)_4(\text{SO}_4)_2]\text{Br}$   
(d)  $[\text{Co}(\text{NH}_3)_4\text{Br}(\text{SO}_4)]$

### SECTION - II

#### More than One Options Correct Type

11. For the given aqueous reactions which of the following statements are true?



- (a) The first reaction is a redox reaction.  
(b) White precipitate is  $\text{Zn}_3[\text{Fe}(\text{CN})_6]_2$ .  
(c) Addition of filtrate to starch solution gives blue colour.  
(d) White precipitate is soluble in  $\text{NaOH}$  solution.
12. The type of isomerism present in nitropentaammine chromium(II) chloride is  
(a) optical (b) linkage  
(c) ionisation (d) polymerisation.
13. For complex ion,  $[\text{Cu}(\text{NH}_3)_4]^{2+}$   
(a) stability constant is  $\frac{[\text{Cu}(\text{NH}_3)_4]^{2+}}{[\text{Cu}^{2+}][\text{NH}_3]^4}$   
(b) stability constant is the inverse of instability constant

- (c) greater is the stability constant, stronger is the Cu—NH<sub>3</sub> bond
- (d) the given complex is only produced when excess of NH<sub>3</sub> solution is added to Cu<sup>2+</sup> solution.

### SECTION - III

#### Paragraph Type

#### Paragraph for Questions 14 and 15

**Exp. (A) :** When Fe(CN)<sub>2</sub> solution is treated with KCN solution, species formed no longer gives tests of Fe<sup>2+</sup> and CN<sup>-</sup>.

**Exp. (B) :** When K<sub>2</sub>SO<sub>4</sub> solution is treated with Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> solution, species formed gives tests of K<sup>+</sup>, Al<sup>3+</sup> and SO<sub>4</sub><sup>2-</sup>.

14. Species formed in experiment (A) does not give test of Fe<sup>2+</sup> and CN<sup>-</sup>. It is due to formation of
- (a) K<sub>2</sub>[Fe(CN)<sub>4</sub>] (b) K<sub>3</sub>[Fe(CN)<sub>5</sub>]  
 (c) K<sub>3</sub>[Fe(CN)<sub>6</sub>] (d) K<sub>4</sub>[Fe(CN)<sub>6</sub>]
15. Species formed in experiment (B) is
- (a) complex (b) double salt  
 (c) liquid crystal (d) none of these.

#### Paragraph for Questions 16 and 17

Double salts are addition compounds which lose their identity in aqueous solution whereas complexes which are also addition compounds do not lose their identity in aqueous solution. The coordination compounds show isomerism and find applications in photography, qualitative analysis, metallurgy, water purification and in the treatment of various diseases.

16. Which of the following statements is incorrect?
- (a) Alum is a double salt.  
 (b) Calcium salt of EDTA is used in the treatment of lead poisoning.  
 (c) Effective atomic number of the metals in complexes [Ni(CO)<sub>4</sub>] and [Fe(CN)<sub>6</sub>]<sup>4-</sup> is same.  
 (d) Tris(triphenylphosphine)rhodium(I) chloride is effective heterogeneous catalyst for hydrogenation of alkenes.
17. Choose the correct option for the complex [PtCl<sub>2</sub>(en)<sub>2</sub>]<sup>2+</sup>.
- (a) Platinum is in +2 oxidation state.  
 (b) Racemic mixture is obtained on mixing mirror images of its *trans*-form in 1 : 1 molar ratio.  
 (c) It has two five membered chelating rings.  
 (d) Both (b) and (c).

### SECTION - IV

#### Matching List Type

18. Match the List I with List II and select the correct answer using the codes given below the lists :

List (Complex)	List II (Hybridisation)		
(P) $[\text{Ni}(\text{CO})_4]$	(i) $sp^3 d^2$		
(Q) $[\text{Ni}(\text{CN})_4]^{2-}$	(ii) $sp^3$		
(R) $[\text{Fe}(\text{CN})_6]^{4-}$	(iii) $d^2 sp^3$		
(S) $[\text{MnF}_6]^{4-}$	(iv) $dsp^2$		
<b>P</b>	<b>Q</b>	<b>R</b>	<b>S</b>
(a) (iv)	(iii)	(ii)	(i)
(b) (ii)	(iv)	(iii)	(i)
(c) (iii)	(iv)	(ii)	(i)
(d) (i)	(ii)	(iii)	(iv)

19. Match the List I with List II and select the correct answer using the codes given below the lists :

List (Complex ion)		List II (Colour)	
(P) $[\text{CoF}_6]^{3-}$		(i) Blue-green	
(Q) $[\text{Co}(\text{NH}_3)_6]^{3+}$		(ii) Pale yellow	
(R) $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$		(iii) Green	
(S) $[\text{Co}(\text{CN})_6]^{3-}$		(iv) Yellow-orange	
P	Q	R	S
(a) (iii)	(iv)	(i)	(ii)
(b) (iii)	(iv)	(ii)	(i)
(c) (i)	(iii)	(iv)	(ii)
(d) (iv)	(i)	(iii)	(ii)

### SECTION - V

#### Assertion Reason Type

20. **Assertion :** The complex ion *trans*-[Co(en)<sub>2</sub>Cl<sub>2</sub>]<sup>+</sup> is optically active.  
**Reason:** It is an octahedral complex.
21. **Assertion :** In complex [Cr(NH<sub>3</sub>)<sub>4</sub>BrCl]Cl, the 'spin only' magnetic moment is close to 1.73 B.M.  
**Reason:** Mononuclear complexes of chromium(III) in strong field ligand have three unpaired electrons.
22. **Assertion :** [Ni(en)<sub>3</sub>]Cl<sub>2</sub> has lower stability than [Ni(NH<sub>3</sub>)<sub>6</sub>]Cl<sub>2</sub>.  
**Reason:** In [Ni(en)<sub>3</sub>]Cl<sub>2</sub>, the geometry of Ni is trigonal bipyramidal.

### SECTION - VI

#### Integer Value Correct Type






23. The number of possible geometrical isomers for the complex [Pt(NO<sub>2</sub>)(py)(NH<sub>2</sub>OH)(NH<sub>3</sub>)] is
24. What is the coordination number of the metal ion in the red complex formed in a qualitative analysis of iron(III)?
25. What is the denticity of N(CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)<sub>3</sub>?

## SOLUTIONS


### ALDEHYDES, KETONES AND CARBOXYLIC ACIDS

1. (b)




2. (d):



5. (d): The  $\gamma$  C—O bond is shortest because it has double bond character. The  $\beta$  C—O bond is shorter because it also has some double bond character due to resonance. The  $\alpha$  C—O bond is longest because it is always a single bond.





6. (d): With calculated amount of Grignard reagent, ketones are formed. But excess of Grignard reagent can lead to the formation of tertiary alcohol. Thus, dimethyl cadmium is the best reagent.




8. (c)






9. (c):








10. (a)



Mechanism :




12. (a, b, c, d) : The main reaction of carbonyl compounds is nucleophilic addition reaction. Reactions with ammonia derivatives are also examples of nucleophilic addition reaction.




14. (c) : Due to  $-I$  effect of Cl.

15. (c) :  $\text{BrCH}_2\text{CH}_2\text{COOH}$  has the smallest dissociation constant (*i.e.*, the weakest acid) since Br has less  $-I$ -effect than F and is far from the  $\text{—COOH}$  group.




17. (a) 18. (a) 19. (d)


20. (d) : The semicarbazone formation occurs with more nucleophilic ( $\text{NH}_2$  side).





21. (c) : It is known that esters with  $\alpha$ -hydrogen form carbanion when treated with base. It brings about nucleophilic substitution at the carbonyl group of the ester to yield  $\beta$ -keto ester. This is known as Claisen condensation reaction.  $\text{LiAlH}_4$  reduces esters to alcohols.




23. (2) :



24. (3) : Formic acid and benzoic acid do not contain  $\alpha$ -H atoms so they cannot give Hell—Volhard—Zelinsky reaction.



25. (5) : Compounds having  $\text{CH}_3\text{—C(=O)—}$  or  $\text{CH}_3\text{—CH(OH)—}$  groups give positive iodoform test.





## COORDINATION COMPOUNDS


1. (a): The configuration will be  $t_{2g}^3 e_g^1$   
 $\therefore \text{CFSE} = 3(-0.4) + 1(0.6) = -0.6 \Delta_o$
2. (a):  $\text{CoCl}_3 \cdot 6\text{NH}_3 \rightarrow x\text{Cl}^- \xrightarrow{\text{AgNO}_3} x\text{AgCl} \downarrow$   
Moles of  $\text{AgCl} = \frac{4.32}{143.5}$   
and moles of  $\text{CoCl}_3 \cdot 6\text{NH}_3 = \frac{2.675}{267.5}$   
or  $\frac{4.32}{143.5} = x \cdot \frac{2.675}{267.5}$   
 $\therefore x = 3$   
Hence, the formula of the complex is  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ .
3. (c): Sodium nitroprusside  $\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}]$  is named as sodium pentacyanonitrosylferrate (II). In this compound, NO exists in +1 and Fe exists in +2 oxidation state.
4. (d): Ferrocene is a sandwich complex in which all the five carbon atoms of cyclopentadienyl anion are linked to the metal by pi-bonds.
5. (a): Octahedral complex,  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$  shows ionisation as well as geometrical isomerism.  
Ionisation isomers are:  
 $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$  and  $[\text{Co}(\text{NH}_3)_4\text{BrCl}]\text{Br}$   
Geometrical isomers are:  

$$\left[ \begin{array}{c} \text{H}_3\text{N} \quad \text{Br} \quad \text{Br} \\ \diagdown \quad | \quad / \\ \text{Co} \\ / \quad | \quad \diagdown \\ \text{H}_3\text{N} \quad \text{NH}_3 \end{array} \right]^+$$

cis-form





$$\left[ \begin{array}{c} \text{H}_3\text{N} \quad \text{Br} \quad \text{NH}_3 \\ \diagdown \quad | \quad / \\ \text{Co} \\ / \quad | \quad \diagdown \\ \text{H}_3\text{N} \quad \text{Br} \quad \text{NH}_3 \end{array} \right]^+$$

trans-form
6. (a)
7. (c): Oxidation state of Cr =  $x + 2(-1) = +1$   
 $\Rightarrow x = +3$   
 $\text{Cr}^{3+}$  occupies 3d orbitals and its coordination number in the given complex is 6.
8. (b): If  $P > \Delta_o$ , it favours the high spin complexes and there is no pairing of d-electrons taken place.  
So,  $3d^4$  will be  $t_{2g}^3 e_g^1$  or  $t_{2g}^{111} e_g^1$ .
9. (c): Oxidation state:  $4(+1) + x + 6(-1) = 0$   
 $4 + x - 6 = 0 \Rightarrow x = +2$   
C.N. = 6, configuration =  $t_{2g}^5 e_g^0$   
 $\text{Mn}^{2+} = 3d^5 4s^0$



It has one unpaired electron and hence it is paramagnetic in nature.


$$\mu = \sqrt{1(1+2)} = \sqrt{3} = 1.732 \text{ B.M.}$$





(a) is correct as in reaction (i),  $\text{I}^- (-1)$  is being oxidised to  $\text{I}_2(0)$  and  $\text{Fe}^{3+}$  is being reduced to  $\text{Fe}^{2+}$ .

(b) is incorrect as white precipitate is of  $\text{K}_2\text{Zn}_3[\text{Fe}(\text{CN})_6]_2$  or  $\text{Zn}_2[\text{Fe}(\text{CN})_6]$ .


(c) is correct as



(d) is correct as white precipitate of  $\text{K}_2\text{Zn}_3[\text{Fe}(\text{CN})_6]_2$  or  $\text{Zn}_2[\text{Fe}(\text{CN})_6]$  is soluble in NaOH as



In,  $-\text{NO}_2$ : N is donating site while in  $-\text{ONO}$ , O is donating site. Thus, it can behave as linkage isomer and Cl is present out of the coordination sphere thus it can behave as ionisation isomer as well.



$$\text{Stability constant } (\beta) = \frac{[\text{Cu}(\text{NH}_3)_4]^{2+}}{[\text{Cu}^{2+}] \cdot [\text{NH}_3]^4}$$

$$\text{Stability constant} \propto \frac{1}{\text{Instability constant}}$$





YOUR WAY **CBSE XII**



Series 6

CHAPTERWISE PRACTICE PAPER :

ALDEHYDES, KETONES AND CARBOXYLIC ACIDS | AMINES

Time Allowed : 3 hours

Maximum Marks : 70


GENERAL INSTRUCTIONS

- All questions are compulsory.
- Q. no. 1 to 5 are very short answer questions and carry 1 mark each.
- Q. no. 6 to 10 are short answer questions and carry 2 marks each.
- Q. no. 11 to 22 are also short answer questions and carry 3 marks each.
- Q. no. 23 is a value based question and carry 4 marks.
- Q. no. 24 to 26 are long answer questions and carry 5 marks each.
- Use log tables if necessary, use of calculators is not allowed.

1. Arrange the following in the decreasing order of ease of hydrolysis:




2. Write down the reactions of ethyl cyanide with




3. Name the alkene which on reductive ozonolysis gives only acetone.

4. Why does aniline dissolve in aqueous  $\text{HCl}$ ?



5. Arrange the following in decreasing order of their acidic strength :



6. Compare the basicity of the following compound:



7. How will you obtain the following?




OR

Give a chemical test to distinguish between

- Formic acid and acetic acid
- Oxalic acid and acetic acid.

8. Complete the following reaction :




9. Write the IUPAC name of the following compounds :


- $(\text{CH}_3)_2\text{CH}-\text{CO}-\text{CH}(\text{CH}_3)_2$
- $\text{Ph}-\text{CH}=\text{CH}-\text{CHO}$

10. Answer the following questions :

- How will you convert an alkyl halide into a primary amine having one carbon atom more than the alkyl halide?
- How will you convert a carboxylic acid into an amine having one carbon atom less than the carboxylic acid?

11. Identify A, B, C and D in the reactions given below :





12. How will you convert the following :

- (i) Nitrobenzene into aniline
- (ii) Primary amine to secondary amine
- (iii) Aniline into *N*-phenylethanamide?

13. Can Gatterman-Koch reaction be considered similar to Friedel-Crafts acylation? Discuss.


14. State the reason for the following :

- (i) Acetylation of aniline reduces its activation effect.
- (ii) Methyl amine is more basic than aniline.
- (iii) Although  $-\text{NH}_2$  is *o/p* directing group, yet aniline on nitration gives a significant amount of *m*-nitroaniline.

15. How will you obtain :

- (i) Butanoic acid from 1-bromopropane
- (ii) Propanoic acid from ethyne
- (iii) 2-Methylbutanoic acid from butan-2-ol?

16. Give the structures of A, B and C in the following reactions :



17. What happens when

- (i) Acetamide is heated with bromine and caustic potash
- (ii) Acetyl chloride reacts with water
- (iii) Acetamide reacts with phosphorus pentoxide?




18. Answer the following questions :

- (i) Why electrophilic substitution in aromatic amines takes place more readily than benzene?
- (ii) Why primary amines have higher boiling point than corresponding tertiary amines?
- (iii) Write two important uses of *N,N*-dimethyl amines (DMA).

OR

- (a) Arrange the following in increasing order of basic strength : Aniline, *p*-nitroaniline and *p*-toluidine.
- (b) State reason for the following :
  - (i) Ethylamine is soluble in water whereas aniline is not.
  - (ii) It is difficult to prepare pure amines by ammonolysis of alkyl halides.

19. Predict the products of the following reactions :



20. Write short notes on the following :

- (i) Carbylamine reaction
- (ii) Diazotisation
- (iii) Gabriel phthalimide synthesis.


21. An organic compound A contains 69.77% carbon, 11.63% hydrogen and rest oxygen. The molecular mass of the compound is 86. It does not reduce Tollens' reagent but forms an addition compound with sodium hydrogen sulphite and gives positive iodoform test. On vigorous oxidation, it gives ethanoic acid and propanoic acids. Derive the structure of compound A.

22. An aromatic compound 'A' on treatment with aqueous ammonia and on heating forms compound 'B' which on heating with  $\text{Br}_2$  and KOH forms a compound 'C' of molecular formula  $\text{C}_6\text{H}_7\text{N}$ . Write the structure and IUPAC name of compounds A, B and C.

23. Sushil's friends want to play Holi with synthetic colours, eggs, muddy water etc. Sunil persuades his friends to play Holi with natural colours. He reminds that last time one of his friend had suffered from skin allergy after playing Holi with synthetic colours. It took long time to recover. Sushil's friends agreed and prepared natural colours using leaves and flowers.

- (i) What is diazotisation reaction?
- (ii) Name two dyes prepared by diazotisation.
- (iii) Write equation for the preparation of any one of the above dyes.
- (iv) What are the values expressed by Sushil?


24. (i) Identify A, B and C in the following sequence of the reactions :




- (ii) Predict the structure of products formed when benzaldehyde is treated with
  - (a) Conc. NaOH
  - (b)  $\text{HNO}_3/\text{H}_2\text{SO}_4$  (at 273-383 K).

(i) Give a chemical test to distinguish between the following :

- (ii) Complete the following reaction :



- (iii)** Identify  $A$ ,  $B$  and  $C$  in the following sequence:



- (i) Benzyl chloride to 2-phenylethanamine
- (ii) Toluene to *o*-chlorotoluene
- (iii) Toluene to *m*-chlorotoluene?

$$\text{C}_5\text{H}_{13}\text{N} \xrightarrow{\text{aq. NaNO}_2/\text{HCl}} \text{Y}$$

(X) Tertiary  
Optically active alcohol

- (i) Identify (X) and (Y). (ii) Is Y optically active?  
(iii) Give structures of intermediate(s), if any, in the formation of Y from X.



26. (i) An organic compound with molecular formula  $C_9H_{10}O$  forms 2, 4-DNP derivative, reduces Tollens' reagent and undergoes Cannizzaro reaction. On vigorous oxidation, it gives 1, 2-benzenedicarboxylic acid. Identify the compound.

- (ii) A compound (A)  $C_5H_8O_2$  liberates  $CO_2$  on reaction with sodium bicarbonate. It exists in two forms neither of which is optically active. It yields (B)  $C_5H_{10}O_2$  on hydrogenation. Compound (B) can be separated into two enantiomers. Write structural formulae for (A) and (B) with reason.


An organic compound (A) of molecular weight 140.5 has 68.32% carbon, 6.4% hydrogen and 25.26% chlorine. Hydrolysis of (A) with dilute acid gives compound (B)  $C_8H_{10}O$ . Compound (B) can be oxidised under mild conditions to form compound (C),  $C_8H_8O$ . Compound (C) forms a phenylhydrazone (D) with phenylhydrazine and gives positive iodoform test. Deduce the structures of (A) to (D) with proper reasoning.

1.  $\text{CH}_3\text{COCl} > (\text{CH}_3\text{CO})_2\text{O} > \text{CH}_3\text{COOC}_2\text{H}_5 > \text{CH}_3\text{CONH}_2$

2. (i) 
$$\text{C}_2\text{H}_5\text{CN} + \text{H}_2\text{O} \xrightarrow[\text{conc. HCl}]{\text{H}_2\text{O}_2(\text{alk.})} \text{C}_2\text{H}_5\text{CONH}_2$$
  
Propanamide





4. Aniline dissolves in aqueous HCl due to the formation of water soluble salt.




5.  $\text{FCH}_2\text{COOH} > \text{ClCH}_2\text{COOH} > \text{C}_6\text{H}_5\text{CH}_2\text{COOH} > \text{CH}_3\text{COOH} > \text{CH}_3\text{CH}_2\text{OH}$


6. The significant difference among these three bases depends on hybridisation of  $\beta$ -carbon atom. The compounds containing carbon with more  $s$ -character show more electron withdrawing nature and hence least basic. Thus, decreasing order of basicity is, (ii) > (i) > (iii).





(i) Formic acid produces silver mirror with ammoniacal silver nitrate solution. Acetic acid does not give this test.




Formic acid      Tollens' reagent



- (ii) Oxalic acid on heating evolve  $\text{CO}_2$  gas. The gas when passed through lime water forms a white precipitate.











Acetic acid does not give this test.






9. (i) 2, 4-Dimethylpent-3-one  
(ii) 3-Phenylprop-2-enal





13. Yes, Friedel-Crafts acylation involves attack of acylium ion,  $(\text{R}-\text{C}^+=\text{O})$  an electrophile on aromatic substrates to give rise to aromatic ketones. Gatterman-Koch reaction involves attack of  $\text{H}-\text{C}\equiv\text{O}^+$  (an electrophile) on aromatic substrates to give rise to aromatic aldehydes.

Friedel - Crafts reaction :








Gatterman-Koch reaction :



Thus, Gatterman-Koch aldehyde synthesis can be considered similar to Friedel-Crafts acylation.

14. (i) After acetylation of aniline, acetanilide is formed in which electron density on N-atom decreases due to the presence of  $-\text{COCH}_3$  group, having  $-I$  effect, and hence, activation effect of aniline gets reduced.
- (ii) In aniline, lone pair of electrons on nitrogen atom is not free for donation because it is involved in resonance. But in case of methylamine, lone pair of electrons on nitrogen atom is free for donation. So, aniline is less basic than methylamine.
- (iii) Nitration is carried out with conc.  $\text{HNO}_3 / \text{H}_2\text{SO}_4$ . In presence of acid, the  $-\text{NH}_2$  group of aniline gets protonated and is converted into  $-\text{NH}_3^+$  group. This positively charged group acts as a strong electron withdrawing and act as a *meta*-directing group. Hence, the incoming electrophile goes to *m*-position and therefore, significant amount of *m*-nitroaniline is formed.





18. (i)  $-\text{NH}_2$  group is electron releasing group, having +R-effect thus increases electron density in benzene ring. So, aniline is more reactive towards electrophilic substitution reaction than benzene.

(ii) Boiling point of amines is high due to presence of hydrogen bonding between amine molecules. In primary amine, large number of hydrogen atoms are available for hydrogen bonding than as compare to tertiary amines. Thus, have higher boiling point.

(iii) (a) It is used as an intermediate in the manufacture of dyes and other substances.

(b) It is important precursor to dyes such as crystal violet.

OR

(a) Electron withdrawing group ( $-\text{NO}_2$ ) on benzene ring decreases the basicity and electron donating group ( $-\text{CH}_3$ ) on benzene ring increases the basicity of the compound.




Increasing order of basic strength :

$p$ -nitroaniline < aniline <  $p$ -toluidine

(b) (i) Ethylamine is soluble in water due to formation of intermolecular hydrogen bonding with water molecules. However, in aniline, due to presence of large hydrophobic aryl group, the extent of hydrogen bonding decreases considerably and hence, insoluble in water.

(ii) The ammonolysis of alkyl halides with ammonia is a nucleophilic substitution reaction

in which a mixture of products is formed and it is not possible to separate individual amines from the mixture.



20. (i) **Carbylamine reaction** : Both aliphatic or aromatic  $1^\circ$  amines on warming with  $\text{CHCl}_3$  in presence of alcoholic KOH solution produce an offensive smell of carbylamines.

(ii) **Diazotisation** : Aromatic diazonium salts are prepared by adding a cold aqueous solution of  $\text{NaNO}_2$  to the solution of an aromatic  $1^\circ$  amine in a mineral acid ( $\text{HCl}$ ,  $\text{H}_2\text{SO}_4$ , etc.) at 273-278 K. This reaction is called diazotisation.

(iii) **Gabriel phthalimide synthesis** : In this reaction, phthalimide is converted into its potassium salt by treating it with alcoholic potassium hydroxide. Further, potassium phthalimide is heated with an alkyl halide to yield an  $N$ -alkylphthalimide which on alkaline hydrolysis gives phthalic acid and primary amine.

21. C = 69.77%, H = 11.63%

$\therefore \text{O} = 100 - (69.77 + 11.63) = 18.6\%$

Element	Percentage	Molar mass	Relative number ratio	Simple ratio
C	69.77	12	5.81	5
H	11.63	1	11.63	10
O	18.6	16	1.16	1

Empirical formula of given compound =  $\text{C}_5\text{H}_{10}\text{O}$

Empirical formula mass =  $5 \times 12 + 10 \times 1 + 1 \times 16 = 86$

$$\Rightarrow n = \frac{86}{86} = 1$$

Since, it does not give Tollens' test but gives positive iodoform test, so it is a methyl ketone derivative *i.e.*, have  $\text{—COCH}_3$  group.

$$\text{CH}_3-\underset{\text{O}}{\underset{\parallel}{\text{C}}}-\text{CH}_2\text{CH}_2\text{CH}_3 \xrightarrow{[\text{O}]} \text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{COOH}$$


Pentan-2-one(A)                      Ethanoic acid                      Propanoic acid


- $$\text{C}_6\text{H}_5\text{CONH}_2 \xrightarrow[\text{(Hoffmann bromamide reaction)}]{\text{Br}_2 + \text{KOH}} \text{C}_6\text{H}_5\text{NH}_2$$
- (B)
(C)

$$\text{C}_6\text{H}_5\text{COOH} \xrightarrow[\text{(ii) } \Delta]{\text{(i) aq. NH}_3} \text{C}_6\text{H}_5\text{CONH}_2$$

(A)
(B)


- (ii) *p*-Hydroxyazobenzene (Orange dye)  
*p*-Aminoazobenzene (Yellow dye)




- (ii) (a) 



- $$\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5 + \text{NaHCO}_3 \longrightarrow \text{No reaction}$$

- (ii) 





# MPP-7 MONTHLY Practice Problems

Class XII

This specially designed column enables students to self analyse their extent of understanding of specified chapter. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.


## Alcohols, Phenols and Ethers

Total Marks : 120

Time Taken : 60 Min.


NEET / AIIMS

Only One Option Correct Type



In the above reaction, the leaving group is

- (a)  $[\text{HOZnCl}_2]^-$  (b)  $\text{H}_2\text{O}$   
(c)  $\text{HO}^-$  (d)  $\text{H}_3\text{O}^+$




In the above reaction sequence, Z is

- (a) toluene (b) cresol  
(c) benzene (d) benzol.

3. In the manufacture of ethanol from sugar the enzymes used are


- (a) diastase and zymase  
(b) maltase and zymase  
(c) diastase and invertase  
(d) invertase and zymase.

4. Which of the following reagents will convert acetophenone to the given alcohol?



- (a)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$  followed by hydrolysis  
(b)  $\text{CH}_3\text{CH}(\text{Br})\text{CH}_3$ ,  $\text{AlCl}_3$   
(c)  $(\text{CH}_3)_2\text{CHMgBr}$  followed by acid hydrolysis  
(d)  $\text{CH}_3\text{CHOHCH}_3$ ,  $\text{Zn}$



5. Which of the following is not expected to be the intermediate of the given reaction?



6. Between *p*-nitrophenol and *p*-cresol, solubility in base is

- (a) almost nil in both the cases  
(b) higher for *p*-nitrophenol  
(c) higher for *p*-cresol  
(d) equal in both the cases.


7. Which of the following sets of compounds cannot turn clear orange solution of  $\text{CrO}_3/\text{dil. H}_2\text{SO}_4$  to greenish blue solution?




- (a) I, IV (b) II, III  
(c) I, II (d) III, IV



8. *p*-Cresol reacts with chloroform in alkaline medium to give a compound A which adds hydrogen cyanide to form compound B. The latter, on acidic hydrolysis gives chiral carboxylic acid. The structure of the carboxylic acid is




9. In the following reaction,



- (a) ethane (b) ethylene  
(c) butane (d) propane.
10. In Williamson's synthesis, ethoxyethane is prepared by
- (a) passing ethanol over heated alumina  
(b) heating sodium ethoxide with ethyl bromide  
(c) treating ethyl alcohol with excess of  $\text{H}_2\text{SO}_4$  at 430–440 K  
(d) heating ethanol with dry  $\text{Ag}_2\text{O}$ .
11. Aluminium *iso*-propoxide is used as a specific reagent for which of the following conversions?
- (a)  $\text{R}-\text{NO}_2 \longrightarrow \text{R}-\text{NOH}$   
(b)  $\text{RCOOR} \longrightarrow \text{RCH}_2\text{OH}$   
(c)  $\text{RCH}=\text{CHCHO} \longrightarrow \text{RCH}=\text{CH}-\text{CH}_2\text{OH}$   
(d)  $\text{RCH}=\text{CH}-\text{CH}_2\text{Cl} \longrightarrow \text{RCH}=\text{CHCH}_3$

12. In which of the following maximum dehydration takes place?



#### Assertion & Reason Type

**Directions :** In the following questions, a statement of assertion is followed by a statement of reason. Mark the correct choice as :

- (a) If both assertion and reason are true and reason is the correct explanation of assertion.  
(b) If both assertion and reason are true but reason is not the correct explanation of assertion.  
(c) If assertion is true but reason is false.  
(d) If both assertion and reason are false.

13. **Assertion :** Anisole undergoes electrophilic substitution at *o*- and *p*-positions.

**Reason :** Anisole is less reactive than phenol towards electrophilic substitution reactions.

14. **Assertion :** Phenols and alcohols can be distinguished by NaOH.

**Reason :** Alcohols are very weak acids as compared to phenols.


15. **Assertion :** Solubility of *n*-alcohols in water decreases with increase in molecular mass.

**Reason :** The proportion of the hydrocarbon part in alcohols increases with increase in molecular mass which permits enhanced hydrogen bonding with water.

#### JEE MAIN / JEE ADVANCED


##### Only One Option Correct Type

16. What is/are the product(s) of the following reaction?




- (a) Only 1  
 (b) 1 : 1 mixture of 2 and 3  
 (c) Only 2  
 (d) 1 : 1 : 1 mixture of 1, 2 and 3


17. An organic compound A reacts with methyl magnesium iodide to form an addition product which on hydrolysis forms the compound B. Compound B gives blue colour salt in Victor Meyer's test. The compounds A and B are respectively  
 (a) acetaldehyde, tertiary butyl alcohol  
 (b) acetaldehyde, ethyl alcohol  
 (c) acetaldehyde, isopropyl alcohol  
 (d) acetone, isopropyl alcohol.
18. Which of the following reactions does not yield phenol?



19. Consider the following reaction,



Low boiling fraction 'Y' is




**More than One Options Correct Type**

20. Which of the following statements about ethers are correct?  
 (a) Ethers are relatively inert compounds.

- (b) Ethers are weakly acidic.  
 (c) Ethers form oxonium salts.  
 (d) Ethers form stable complexes with Lewis acids.

21. Propan-1-ol and propan-2-ol can be best distinguished by  
 (a) oxidation with alkaline  $\text{KMnO}_4$  followed by reaction with  $\text{H}_2\text{O}$   
 (b) oxidation with PCC followed by reaction with Tollens' reagent  
 (c) heating with copper followed by reaction with iodoform  
 (d) reaction with conc.  $\text{H}_2\text{SO}_4$  followed by reaction with Fehling's solution.

22. Which of the following reactions give 2-butanol as a product?



23. Diethyl ether can be distinguished from *n*-butanol by  
 (a) aqueous  $\text{FeCl}_3$  solution  
 (b) reaction with Na metal  
 (c) Tollens' reagent  
 (d) reaction with chromic anhydride ( $\text{CrO}_3$ ) in dil.  $\text{H}_2\text{SO}_4$ .

### Winners of September 2017 Crossword

- Devjit Acharjee, West Bengal
- Ajay Chatterji, Odisha

### Solution Senders of Chemistry Musing

#### Set - 51

- Sukanya Sahoo, West Bengal
- Aradhana Chopra, New Delhi

#### Set - 50

- Anshuman Pan, West Bengal
- Kanav Kapoor, Uttar Pradesh

### Integer Answer Type

24. Among the following, the number of alcohols showing iodoform test is  
 $\text{CH}_3\text{OH}$ ,  $\text{CH}_3\text{CH}_2\text{OH}$ ,  $(\text{CH}_3)_2\text{CHOH}$ ,  
 $(\text{CH}_3)_3\text{C}-\text{OH}$ ,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ ,  
 $\text{CH}_3\text{CHOHCH}_2\text{CH}_3$ ,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ ,  
 $(\text{C}_2\text{H}_5)_2\text{CHOH}$ ,  $\text{CH}_3\text{CHOHCH}(\text{CH}_3)_2$
25. How many of the following substances are less acidic than ethyl alcohol?  
 Water, acetylene, carbonic acid, methanol, *iso*-propyl alcohol, phenol, *tert*-butyl alcohol, *n*-propyl alcohol
26. How many ethers will be formed when a mixture of ethyl alcohol and methyl alcohol is treated with conc.  $\text{H}_2\text{SO}_4$ ?

### Comprehension Type

The cleavage of ethers to give the original alkyl halide and the alcohol is carried out by heating the ether with a halogen acid. Usually, HI at 373 K is used, but some ethers can be cleaved even by HBr or HCl. Depending upon the nature of the alkyl/aryl groups around the oxygen atom and the polarity of the solvent in which HI is dissolved, the reaction may follow either  $\text{S}_{\text{N}}2$  or  $\text{S}_{\text{N}}1$  mechanism.

27. Which of the following ethers is cleaved even by hydrogen chloride at room temperature?  
 (a)  $\text{C}_6\text{H}_5-\text{O}-\text{CH}_2\text{CH}_3$   
 (b)  $\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_3$   
 (c)  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_2\text{CH}_3$   
 (d)  $(\text{CH}_3)_3\text{C}-\text{O}-\text{C}(\text{CH}_3)_3$
28. The major products formed when the following ether is heated with conc. HI  
 $\text{CH}_3\text{CH}_2\text{CH}_2-\text{O}-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$   
 are  
 (a)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{I} + \text{HO}-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$   
 (b)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + \text{I}-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$   
 (c)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{I} + \text{I}-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$   
 (d)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + (\text{CH}_3)_2\text{C}=\text{CHCH}_3$

### Matrix Match Type

29. Match column I with column II and choose the correct option using the codes given below :

Column I	Column II
(A) Obtained by reaction between ethyl acetate and excess of $\text{CH}_3\text{MgBr}$ followed by acidic hydrolysis	(p) $\text{CH}_3\text{CH}_2\text{OH}$
(B) Gets easily oxidised by $\text{K}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4$	(q) $\text{CH}_3\text{CH}_2\text{CHOHCH}_3$
(C) Produces blue colouration in Victor-Meyer's test	(r) $\text{C}_6\text{H}_5\text{OH}$
(D) Produces violet colouration with neutral $\text{FeCl}_3$ solution	(s) $(\text{CH}_3)_3\text{COH}$

A	B	C	D
(a) p	s	r	q
(b) s	p	q	r
(c) q	r	p	s
(d) r	q	s	p

30. Match column I with column II and choose the correct option using the codes given below :

Column I	Column II
(A) Phenol + Neutral $\text{FeCl}_3$	(p) No reaction
(B) Phenol + $\text{Br}_{2(\text{aq})}$	(q) Violet colour
(C) Phenol + $\text{NaHCO}_3$	(r) White ppt.
(D) Picric acid + $\text{NaHCO}_3$	(s) $\text{CO}_2$ gas is evolved

A	B	C	D
(a) p	r	q	s
(b) s	p	q	r
(c) q	r	p	s
(d) p	q	s	r



Keys are published in this issue. Search now! ☺

## SELF CHECK



### Check your score! If your score is

No. of questions attempted	.....	> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
No. of questions correct	.....	90-75%	GOOD WORK !	You can score good in the final exam.
Marks scored in percentage	.....	74-60%	SATISFACTORY !	You need to score more next time.
		< 60%	NOT SATISFACTORY !	Revise thoroughly and strengthen your concepts.


# CHEMISTRY MUSING

## PROBLEM SET 52

**C**hemistry Musing was started from August '13 issue of Chemistry Today. The aim of Chemistry Musing is to augment the chances of bright students preparing for JEE (Main and Advanced) / NEET / AIIMS / JIPMER with additional study material. In every issue of Chemistry Today, 10 challenging problems are proposed in various topics of JEE (Main and Advanced) / NEET. The detailed solutions of these problems will be published in next issue of Chemistry Today. The readers who have solved five or more problems may send their solutions. The names of those who send at least five correct solutions will be published in the next issue. We hope that our readers will enrich their problem solving skills through "Chemistry Musing" and stand in better stead while facing the competitive exams.

### JEE MAIN/NEET

- The pyknometric density of NaCl crystal is  $2.165 \times 10^3 \text{ kg m}^{-3}$  while its X-ray density is  $2.178 \times 10^3 \text{ kg m}^{-3}$ . The fraction of unoccupied sites in NaCl crystal is  
(a) 5.96 (b)  $5.96 \times 10^{-2}$   
(c)  $5.96 \times 10^{-1}$  (d)  $5.96 \times 10^{-3}$
- 'A' on heating at higher temperature ( $>700^\circ\text{C}$ ) in air gives a white infusible amorphous powder B which is decomposed when heated in the current of steam to give white powder 'C' and a gas 'D'. 'D' turns red litmus blue and in aqueous solution, gives reddish brown ppt. with  $\text{K}_2\text{HgI}_4$ . Compound 'C' on strong heating gives 'E'. Here, E is  
(a)  $\text{B}_2\text{O}_3$  (b)  $\text{SiO}_2$  (c)  $\text{NH}_3$  (d)  $\text{N}_2\text{O}$
- Hexavalent uranium mineral such as carnotite is leached with soda ash, the dissolution of uranium takes place by forming uranyl tricarbonate ion. Leaching reaction of uranium is accompanied by generation of  $\text{OH}^-$ . The generation of  $\text{OH}^-$  ion makes the slurry more alkaline and uranium gets precipitated in the form of  
(a)  $\text{Na}_2\text{UO}_4$  (b)  $\text{Na}_2\text{U}_3\text{O}_7$   
(c)  $\text{Na}_4(\text{UO}_2)(\text{CO}_3)_3$  (d)  $\text{UO}_2\text{SO}_4$
- Ellingham diagram for some oxides of P, Q, R, S, T and U is given.



Select the incorrect statement.

- Below  $400^\circ\text{C}$  oxide of P should decompose into metal and  $\text{O}_2$ .
  - Extraction of A is possible by reduction of its oxide by T and U.
  - At  $500^\circ\text{C}$  reduction of oxide of P, Q is possible by R.
  - Reduction of  $\text{T}_2\text{O}_3$  to T is possible by R below  $1000^\circ\text{C}$ .
5. For the equilibrium,  
 $\text{LiCl} \cdot 3\text{NH}_3(\text{s}) \rightleftharpoons \text{LiCl} \cdot \text{NH}_3(\text{s}) + 2\text{NH}_3(\text{g})$   
 $K_p = 9 \text{ atm}^2$ , at  $40^\circ\text{C}$ . A 5 litre flask contains 0.1 mole of  $\text{LiCl} \cdot \text{NH}_3$ . How many moles of  $\text{NH}_3$  should be added to the flask at this temperature to drive the backward reaction for completion?  
 (a) 0.58 (b) 0.2 (c) 0.78 (d) 0.4


### JEE ADVANCED


6. The number of unpaired electrons in octahedral complexes of  $d^9$  and  $d^4$  ions when  $\Delta_o > P$  and  $\Delta_o < P$ .
- |     | $d^9(\Delta_o > P)$ | $d^9(\Delta_o < P)$ | $d^4(\Delta_o > P)$ | $d^4(\Delta_o < P)$ |
|-----|---------------------|---------------------|---------------------|---------------------|
| (a) | 1                   | 1                   | 4                   | 4                   |
| (b) | 1                   | 3                   | 2                   | 4                   |
| (c) | 1                   | 1                   | 2                   | 4                   |
| (d) | 1                   | 0                   | 4                   | 4                   |

### COMPREHENSION


The word "aromatic" though started with benzene and its derivatives only, now it signifies a large variety of organic compounds. The compound is said to be aromatic if it is cyclic planar and contain  $(4n + 2)\pi$  electrons. When the closed loop contains  $4n$  electrons, the system is rather less stable or anti-aromatic.

7. Which of the following has the highest value of dipole moment?






8. Which of the following is the most stable isomer of the hydrocarbon,  $C_{10}H_{10}$ ?



#### INTEGER VALUE

9. The substituents  $R_1$  and  $R_2$  for nine peptides are listed in the table given below. How many of these peptides are positively charged at pH 7.0?



Peptide	$R_1$	$R_2$
I	H	H
II	H	$\text{CH}_3$
III	$\text{CH}_2\text{COOH}$	H
IV	$\text{CH}_2\text{CONH}_2$	$(\text{CH}_2)_4\text{NH}_2$
V	$\text{CH}_2\text{CONH}_2$	$\text{CH}_2\text{CONH}_2$
VI	$(\text{CH}_2)_4\text{NH}_2$	$(\text{CH}_2)_4\text{NH}_2$
VII	$\text{CH}_2\text{COOH}$	$\text{CH}_2\text{CONH}_2$
VIII	$\text{CH}_2\text{OH}$	$(\text{CH}_2)_4\text{NH}_2$
IX	$(\text{CH}_2)_4\text{NH}_2$	$\text{CH}_3$

10. For estimating ozone in the air, a certain volume of air is passed through an alkaline KI solution when  $\text{O}_2$  is evolved and iodide is oxidised to iodine. When such a solution is acidified, the free iodine is evolved which can be titrated with standard  $\text{Na}_2\text{S}_2\text{O}_3$  solution. In an experiment, 9 L of air at 1 atm and  $27^\circ\text{C}$  were passed through an alkaline KI solution, and at the end, the iodine was entrapped in a solution which on titration required 1.5 mL of 0.01 N  $\text{Na}_2\text{S}_2\text{O}_3$  solution. If the volume percentage of ozone in the sample is  $x \times 10^{-3}$  then  $x$  is

**ATTENTION COACHING INSTITUTES:**  
a great offer from MTG

**MTG** offers "Classroom Study Material" for JEE (Main & Advanced), NEET and FOUNDATION MATERIAL for Class 6, 7, 8, 9, 10, 11 & 12 **with YOUR BRAND NAME & COVER DESIGN.**

This study material will save you lots of money spent on teachers, typing, proof-reading and printing. Also, you will save enormous time. Normally, a good study material takes 2 years to develop. But you can have the material printed with your logo delivered at your doorstep.

Profit from associating with MTG Brand — the most popular name in educational publishing for JEE (Main & Advanced)/NEET/PMT....

Order sample chapters on Phone/Fax/e-mail.


Phone : 0124-6601200

09312680856

e-mail : sales@mtg.in | www.mtg.in



#### CLASSROOM STUDY MATERIAL



Your logo here






# Chemistry

## Nobel Prize Winners

### 2017



**Jacques Dubochet**

**Born :** 1942, Aigle, Switzerland

**Affiliation at the time of the award :**  
University of Lausanne, Lausanne,  
Switzerland



**Joachim Frank**

**Born:** 1940, Siegen, Germany

**Affiliation at the time of the award :**  
Columbia University, New York, USA



**Richard Henderson**

**Born:** 1945, Edinburgh, Scotland

**Affiliation at the time of the award :**  
MRC Laboratory of Molecular Biology,  
Cambridge, United Kingdom

The Nobel prize in Chemistry 2017 was awarded to Jacques Dubochet, Joachim Frank and Richard Henderson for the development of cryo-electron microscopy, which simplifies and improves the imaging of biomolecules. This method has moved biochemistry into a new era.

## Molecules of Life, Captured in 3D

Scientific breakthroughs often build upon the successful visualisation of objects invisible to the human eye. However, biochemical maps have long been filled with blank spaces because the available technology has difficulty in generating images of life's molecular machinery. Cryo-electron microscopy changes all of this. Researchers can now freeze biomolecules mid-movement and visualise processes they have never previously seen, which is decisive for both the basic understanding of life's chemistry and for the development of pharmaceuticals. Electron microscopes were long believed to only be suitable for imaging dead matter, because the powerful electron beam destroys biological material. But in 1990, Richard Henderson succeeded in using an electron microscope to generate a three-dimensional image of a protein at atomic resolution. This breakthrough proved the technology's potential.

Joachim Frank made the technology generally applicable. Between 1975 and 1986, he developed an image processing method in which the electron microscope's fuzzy two-dimensional images are analysed and merged to reveal a sharp three-dimensional structure.

Jacques Dubochet added water to electron microscopy. Water evaporates in the electron microscope's vacuum, which makes the biomolecules collapse. In the early 1980s, Dubochet succeeded in vitrifying water – he cooled water so rapidly that it solidified in its liquid form around a biological sample, allowing the biomolecules to retain their natural shape even in a vacuum.

Following these discoveries, the electron microscope's every nut and bolt have been optimised. The desired atomic resolution was reached in 2013, and researchers can now routinely produce three-dimensional structures of biomolecules. In the past few years, scientific literature has been filled with images of everything from proteins that cause antibiotic resistance, to the surface of the Zika virus. Biochemistry is now facing an explosive development and is all set for an exciting future.



# CONCEPT BOOSTER

Hello My family!! Wish you all enjoyed a lot in the festive season and side by side definitely have studied to keep the brain engine smooth and energetic. This is the time when you need a proper planning how to proceed in the forthcoming months with your preparation. Be convinced you will get me always beside you. I will start giving study plan and sequence of study in the forthcoming issues along with some off topics which you need to study. In the beginning, in this issue I have discussed JAHN-TELLER EFFECT, a very important but less discussed topic of COORDINATION CHEMISTRY. Keep your eyes intact on CHEMISTRY TODAY CONCEPT BOOSTER always for further guidance.

\*Arunava Sarkar

## SPECIAL TOPIC OF COORDINATION CHEMISTRY

At the beginning, know the fact that Jahn-Teller effect or Jahn-Teller theorem give an idea about the additional stabilisation of the system.


### JAHN-TELLER THEOREM

Any non-linear system in its orbitally degenerate electronic state will be unstable and therefore in order to attain stability this electronic state will undergo distortion in its geometry and thus will cause a splitting in its orbitally degenerate electronic state. This theorem was developed in the year 1937. Remember that orbitally degenerate electronic state is that state which represents more than one electronic arrangement with the same energy.


Jahn-Teller distortion is therefore that phenomena where distortion in the geometry of the non-linear system takes place in order to gain more stability.

Jahn-Teller distortion lowers the symmetry of the system in such a way that it lowers the energy of the system. It also takes place automatically.


The best example to understand Jahn-Teller distortion is  $\text{Cu}^{2+}$ :



With a perfect octahedral geometry, this configuration can be given as:



Now, two options are here:




### OPTION 1

The  $d$ -electron charge density will be higher in  $z$ -direction than in  $x$ - or  $y$ -direction. So, the nuclear charge of  $\text{Cu}^{2+}$  rather the positive charge of  $\text{Cu}^{2+}$  will be more in  $z$ -direction rather in  $x$ - or  $y$ -direction. As a result, the negative charge of the ligands along the  $z$ -direction will be less attractive than that of along  $x$ - or  $y$ -direction.

Now think the scenario! Due to lesser attraction, ligands along  $z$ -direction will move away from the metal ion whereas ligands along  $x$ - and  $y$ -directions will move nearer to the metal ion.


Obviously, now the octahedral geometry will get distorted to tetragonal geometry and it will be with the elongation along  $z$ -directions and compression along  $x$ - and  $y$ -directions.

So, what we found ultimately? There is elongation of metal-ligand bonds along  $z$ -direction. That is why the  $d$ -electrons of the metal ion experience lesser electrostatic repulsion force from the negative charge on the ligands.



(Distortion of the octahedral geometry to tetragonal geometry due to Jahn-Teller effect)

The similar repulsion force will be more felt at the  $x$ - and  $y$ -directions. So, due to all these  $d_{z^2}$  will be of lower energy and  $d_{x^2-y^2}$  will be of higher energy. In this way the degeneracy gets disturbed and gets lifted.




(Lifting of degeneracy of  $e_g$  orbitals due to Jahn-Teller effect when more electrons are there in  $d_{z^2}$  than  $d_{x^2-y^2}$ )

Now, there is a catch here. No external energy supply is required for the splitting here. It is automatic. Hence, overall energy of the split orbitals will be equal to the energy of the unsplit orbitals. So, if we assume that before the splitting, the energy of the orbitals was  $E$ , then after the splitting total energy of  $d_{z^2}$  and  $d_{x^2-y^2}$  will be  $E$ . This is commonly known as 'Centre of gravity rule' according to which the centre of gravity of the split orbitals is maintained.


So, in a nut shell, when there are more electrons in  $d_{z^2}$  rather than in  $d_{x^2-y^2}$  of an octahedral complex of any metal ion, distortion of octahedral geometry to tetragonal geometry will occur by elongation of metal-ligand bonds along  $z$ -direction.

### OPTION 2

Here,  $d_{x^2-y^2}$  has two electrons and  $d_{z^2}$  has one electron. This is why the negative charges on the ligands along  $x$ - and  $y$ - directions are less attracted by the nucleus or rather nuclear charge. This is of course because of the more effective screening. This screening is of course less along the  $z$ -direction as there is only one electron in  $d_{z^2}$ . As a consequence, there is elongation of metal-ligand bond along  $x$ - and  $y$ -directions whereas there is contraction of metal-ligand bond across  $z$ -direction. As a result the octahedral geometry of  $\text{Cu}^{2+}$  ion gets distorted and it takes tetragonal geometry.



Also, electrostatic repulsion experienced by  $d$ -electrons of  $\text{Cu}^{2+}$  from the negative charge on ligands along  $x$ - and  $y$ -direction will be less than the similar force experienced by the  $d$ -electron(s) from the negative charge along the  $z$ -direction. This is why  $d_{x^2-y^2}$  becomes of lower energy and  $d_{z^2}$  becomes of higher energy. So degeneracy of two orbitals get lifted.



Now, we will take a few aspects which must be understood properly.

First of all, remember that apart from the  $t_{2g}^6 e_g^3$  configuration (ground state) resulting into the formation of octahedral complex, other ground state configuration which show Jahn-Teller effect is  $t_{2g}^6 e_g^1$ . The point to be noted here is in both of these configurations  $t_{2g}$  is uniformly filled up and the type of distortion here depends upon  $e_g$  orbital *i.e.* whether the electron occupies  $d_{x^2-y^2}$  or  $d_{z^2}$  orbital

### Exam Update

### JEE (Advanced) 2018

The Joint Entrance Examination (Advanced) 2018 will be conducted by the IITs. The performance of a candidate in this examination will form the basis for admission to the Bachelor's, Integrated Master's and Dual Degree programs (entry at the 10+2 level) in all the IITs. The decisions of the JAB 2018 will be final in all matters related to JEE (Advanced) 2018 and admission to IITs.

Candidates should be among the top 2,24,000 (including all categories) by scoring positive marks in Paper-1 of JEE (Main)-2018.

Examination will be held on May 20, 2018. The entire JEE (Advanced) 2018 Examination will be conducted in fully computer based test mode. The exam consists of two papers, Paper 1 and Paper 2, each of three hours' duration, and will be held in two sessions. Both the papers are compulsory.

For more information visit [www.jeeadv.ac.in](http://www.jeeadv.ac.in)

### MPP-7 CLASS XII

### ANSWER KEY

- |           |           |           |         |             |
|-----------|-----------|-----------|---------|-------------|
| 1. (a)    | 2. (c)    | 3. (d)    | 4. (c)  | 5. (a)      |
| 6. (b)    | 7. (d)    | 8. (c)    | 9. (a)  | 10. (b)     |
| 11. (c)   | 12. (b)   | 13. (b)   | 14. (a) | 15. (c)     |
| 16. (a)   | 17. (c)   | 18. (b)   | 19. (d) | 20. (a,c,d) |
| 21. (b,c) | 22. (a,b) | 23. (b,d) | 24. (4) | 25. (4)     |
| 26. (3)   | 27. (d)   | 28. (b)   | 29. (b) | 30. (c)     |



# ADVANCED CHEMISTRY BLOC

(RE-READING ORGANIC CHEMISTRY)




Mukul C. Ray, Odisha

In a span of six episodes, we would be covering the indispensable points of organic chemistry, exactly what you need to understand or memorise in this column. Things are really simple in organic chemistry. You just take one step, means one concept or one reaction at a time. Rehearse and progress daily, without fail, even though for 30 minutes. Soon you will realise the things come to your mind effortlessly.

## ALIPHATIC HYDROCARBONS

### 1. Halogenation of alkanes


Chlorination of methane




Step-2 is endothermic but step-3 is exothermic, thus overall process is exothermic. For all other alkanes, both steps are exothermic.

In chlorination, the selectivity of  $\text{Cl}\cdot$  towards primary, secondary and tertiary hydrogen is in the ratio of 1 : 3.8 : 5.

This helps us to calculate the product distribution.




Relative amount of *t*-butyl chloride is 5 while for 2-methyl-1-chloropropane it is 9. No doubt, the later one is the major product.



Together there will be 10 isomers of product possible from the above reaction.


For bromination of methane, step-2 is endothermic but step-3 is exothermic. The overall process is weakly exothermic.

The selectivity ratio of  $\text{Br}\cdot$  towards 1°, 2° and 3° is 1 : 82 : 1600. So no need to calculate, you just go for tertiary product as the major product always.



### 2. Nitration of alkanes

Nitration involves C—C bond cleavage along with C—H bond.



### 3. Physical properties of alkanes


Melting point :  $\text{C}_2\text{H}_6 > \text{CH}_4 > \text{C}_3\text{H}_8$

From propane onwards, melting point gradually increases though not regularly. The difference is less between odd and even as compared to that while moving from even to odd in case of *n*-alkanes. This is the “oscillation” or “alteration rule.”

Check the data (No need to remember) :

Alkane	$\text{C}_3\text{H}_8$	<i>n</i> - $\text{C}_4\text{H}_{10}$	<i>n</i> - $\text{C}_5\text{H}_{12}$	<i>n</i> - $\text{C}_6\text{H}_{14}$
M.P.	-189.9 °C	-135 °C	-131.5 °C	-94.3 °C


And order of melting point :




Boiling points for *n*-alkanes show a smooth increase along the series. For identical number of carbon containing compounds boiling point decreases with increase in branching.

#### 4. Preparation of alkenes

Strong base and heating condition on alkyl halides mostly certifies E<sub>2</sub> mechanism. Generally, it gives Saytzeff product except when -F or -NH<sub>3</sub><sup>+</sup> is the leaving group or bulky base is present where Hofmann elimination gives dominating product.




Branched alkyl halides, Ag<sup>+</sup> catalyst, weaker base like H<sub>2</sub>O or CH<sub>3</sub>OH and heating condition confirms E<sub>1</sub> mechanism. Saytzeff product is always the major product.



Zn is a dehalogenating agent that causes anti-elimination.

Remember the following table. Carefully check there is a symmetry.


Alkene	Addition	Product
<i>cis</i>	<i>Syn</i>	<i>Meso</i>
<i>cis</i>	<i>Anti</i>	<i>dl</i> (±)
<i>trans</i>	<i>Anti</i>	<i>Meso</i>
<i>trans</i>	<i>Syn</i>	<i>dl</i> (±)





#### 5. Addition reactions on alkenes

Alkenes generally follow addition electrophilic (AdE) mechanism.

Addition of HX and H<sub>3</sub>O<sup>+</sup> : Alkene adds H<sup>+</sup> ion across double bond to give a carbocation, which rearranges and then pick the nucleophile.



But addition of Br<sub>2</sub>, Cl<sub>2</sub>/H<sub>2</sub>O involve a cyclic halonium ion. Hence, no rearrangement takes place. The net result is anti-addition.





# CHEMISTRY MUSING

## SOLUTION SET 51


$$1. (d): M_{\text{solute}} = \frac{K_f \times w_{\text{solute}} \times 1000}{W_{\text{solvent}} \times \Delta T_f} = \frac{4.90 \times 3.26 \times 1000}{226 \times 0.112} = 631 \text{ g mol}^{-1}$$

Now, molecular mass of  $\text{Se}_x = x \times 79$

$$631 = x \times 79 \Rightarrow x = 8$$

$\therefore$  Molecular formula of selenium =  $\text{Se}_8$

2. (c):



3. (c): Given:  $T_i = 17^\circ\text{C} = 290 \text{ K}$ ;  $T_f = 57^\circ\text{C} = 330 \text{ K}$

$$\pi_i = 580 \text{ torr} = \frac{580}{760} \text{ atm}; \pi_f = 165 \text{ torr} = \frac{165}{760} \text{ atm}$$

Using equation,  $\pi = CRT$



$$\frac{580}{760} = C_i \times R \times 290 \quad \dots(i)$$

$$\frac{165}{760} = C_f \times R \times 330 \quad \dots(ii)$$



Dividing eqn (i) by (ii), we get

$$\frac{C_i}{C_f} = \frac{2}{1/2} = 4 \Rightarrow \text{Dilution} = 4 \text{ times}$$

4. (d): The compound combines with one mole of hydrogen to form normal hexane, hence the hydrocarbon is *n*-hexene.










5. (d): A compound which forms stable carbocation undergoes solvolysis.



Hence, compounds given in options (b) and (c) do not undergo solvolysis.

6. (c): (a)  $\text{Mg}_2\text{C}_3 + 4\text{H}_2\text{O} \longrightarrow 2\text{Mg}(\text{OH})_2 + \text{C}_3\text{H}_{4(g)}$  (X)




7. (c): On increasing concentration of  $\text{NH}_3$ , the concentration of  $\text{H}^+$  ion decreases.


$$E_{\text{red}} = E^\circ_{\text{red}} - \frac{0.0591}{n} \log[\text{H}^+]$$

$$E_{\text{red}} = 0 - \frac{0.0591}{1} \log 10^{-11} = -0.0591 \times (-11) = 0.65$$

8. (c):  $\text{NH}_3$  has no effect on the standard reduction potential.



Conversion of sodium chromate into sodium dichromate.



In  $\text{FeO} \cdot \text{Cr}_2\text{O}_3$  (X); oxidation number of Cr = +3

In  $\text{Na}_2\text{Cr}_2\text{O}_7$  (Z); oxidation number of Cr = +6

10. (1):  $\pi = iCRT$

For 1 M  $\text{Na}_2\text{SO}_4$ ;  $i = 3$ ,  $\pi = 3RT$

For 1 M  $\text{KCl}$ ;  $i = 2$ ,  $\pi = 2RT$

For 1.5 M  $\text{K}_2\text{SO}_4$ ;  $i = 3$ ,  $\pi = 4.5RT$



For 1.5 M  $\text{NaCl}$ ;  $i = 2$ ,  $\pi = 3RT$

For 2 M  $\text{Al}_2(\text{SO}_4)_3$ ;  $i = 5$ ,  $\pi = 10RT$

For 0.5 M  $\text{BaSO}_4$ ;  $i = 2$ ,  $\pi = 1RT$

Hence, only one compound, i.e., 1.5 M  $\text{NaCl}$  has same osmotic pressure as that of 1 M  $\text{Na}_2\text{SO}_4$  solution.  $\diamond \diamond$

# Now, save up to Rs 2,020\*



## Subscribe to MTG magazines today.

Our 2017 offers are here. Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to [www.mtg.in](http://www.mtg.in) now to subscribe online.

\*On cover price of ₹ 30/- each.

For JEE  
(Main & Advanced),  
NEET, AIIMS AND  
JIPMER

## About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines-Physics For You, Chemistry Today, Mathematics Today & Biology Today-ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

Trust of over 1 Crore readers since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team

- comprising experts and members from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement –

- confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET toppers
- **Bonus:** Exposure to competition at a global level, with questions from Intl. Olympiads & Contests

## SUBSCRIPTION FORM

Please accept my subscription to:

Note: Magazines are despatched by Book-Post on 4<sup>th</sup> of every month (each magazine separately).

☒ Tick the appropriate box.

### PCMB combo

☐ 1 yr: ₹ 1,000 (save ₹ 440) ☐ 2 yr: ₹ 1,800 (save ₹ 1,080) ☐ 3 yr: ₹ 2,300 (save ₹ 2,020)

### PCM combo

☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yr: ₹ 1,500 (save ₹ 660) ☐ 3 yr: ₹ 1,900 (save ₹ 1,340)

### PCB combo

☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yr: ₹ 1,500 (save ₹ 660) ☐ 3 yr: ₹ 1,900 (save ₹ 1,340)

### Individual magazines

☐ Physics ☐ Chemistry ☐ Mathematics ☐ Biology

☐ 1 yr: ₹ 330 (save ₹ 30) ☐ 2 yr: ₹ 600 (save ₹ 120) ☐ 3 yr: ₹ 775 (save ₹ 305)

Enclose Demand Draft favouring **MTG Learning Media (P) Ltd.** payable at New Delhi. You can also pay via Money Orders. Mail this Subscription Form to Subscription Dept., **MTG Learning Media (P) Ltd.**, Plot 99, Sector 44, Gurgaon - 122 003 (HR).

Want the magazines by courier; add the courier charges given below:

☐ 1 yr: ₹ 240 ☐ 2 yr: ₹ 450 ☐ 3 yr: ₹ 600

☒ Tick the appropriate box.

☐ Student ☐ Class XI ☐ XII ☐ Teacher ☐ Library ☐ Coaching

Name: \_\_\_\_\_

Complete Postal Address: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



Pin Code       Mobile #

Other Phone # 0

Email: \_\_\_\_\_

E-mail [subscription@mtg.in](mailto:subscription@mtg.in). Visit [www.mtg.in](http://www.mtg.in) to subscribe online. Call (0)8800255334/5 for more info.

# Presenting India's No.1 NEET Guides



MTG's Complete NEET Guides are India's best selling PMT books!! Rich in theoretical knowledge with a vast question bank comprising a wide variety of problems and exercises, these guidebooks ensure students are ready to compete in the toughest of medical entrance tests. 100% NCERT based, the guidebooks have been updated to match the syllabus and the exam pattern for medical entrance exams. No wonder these guidebooks emerged as the bestsellers in a short period of time.

## HIGHLIGHTS:

- 100% NCERT based
- Comprehensive Chapterwise theory complemented with concept maps, flowcharts and easy-to-understand illustrations
- Last 10 years' questions (2008-2017) of AIPMT/NEET
- Chapterwise Topicwise MCQs with detailed explanations and solutions
- NEET 2017 Solved Paper included
- Over 50% of questions that appeared in NEET 2017 were from MTG's Complete NEET Guides



Scan now with your  
smartphone or tablet\*



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or e-mail: info@mtg.in


\*Application to read QR codes required

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!



# THE PERFECT COACH

Now also available for JEE!



Skill. Passion. Hard work and determination. As a student sitting for the highly competitive JEE, you need all that. However, only a few will win, very likely with the help of a champion coach.

MTG's Champion Series for JEE is just the coach you need. It will guide you in identifying what's important for success and what's not. And then help you check your readiness with its most comprehensive question bank. So you know your strengths and weaknesses right from the word go and course-correct accordingly. Put simply, MTG's Champion Series will help you manage your preparation effort for JEE for maximum outcome. The best part is you study at a pace you're comfortable with.




Because it's all chapterwise, topicwise.



Visit [www.MTG.in](http://www.MTG.in) to buy online. Or visit a leading bookseller near you.  
For more information, email [info@mtg.in](mailto:info@mtg.in) or call 1800 300 23355 (toll-free) today.



The most comprehensive **question bank books**  
that you cannot afford to ignore



30 Years' Physics, Chemistry & Biology contain not only chapterwise questions that have appeared over the last 30 years in NEET/AIPMT, but also full solutions, that too by experts. Needless to say, these question banks are essential for any student to compete successfully in NEET.

#### HIGHLIGHTS:

- Chapterwise questions of last 30 years' (2017-1986) of NEET/AIPMT
- Chapterwise segregation of questions to help you assess the level of effort required to succeed
- An unmatched question bank series with close to 1,000 pages having detailed solutions by experts
- Fully solved questions of NEET 2017 included



Scan now with your  
smartphone or tablet\*



Available at all leading book shops throughout India.  
For more information or for help in placing your order:  
Call 0124-6601200 or email [info@mtg.in](mailto:info@mtg.in)


\*Application to read QR codes required

Visit  
[www.mtg.in](http://www.mtg.in)  
for latest offers  
and to buy  
online!




# Study right. Dig deep.


Build a solid foundation for success  
in JEE Main



₹ 750



₹ 850



₹ 750

**Are you a do-it-yourself type of a student?** Then for success in JEE Main, choose MTG's JEE Main combo, comprising coursebooks for Physics, Chemistry & Mathematics. This combo is all class 11 and 12 students need for a solid and deep understanding of concepts in these three key subjects.

## FEATURES:

- Based on latest pattern of JEE Main
- Full of graphic illustrations & MCQs for deep understanding of concepts
- Covers the entire syllabus
- NCERT Drill MCQs framed from NCERT Books
- 11 Years (2017-2007) Previous Years MCQs of JEE Main / AIEEE
- 2017 JEE Main (Offline & Online) Solved Paper included



Scan now with your  
smartphone or tablet  
Application to read  
QR codes required


Note: Coursebooks are also available separately.

Available at all leading book shops throughout India. To buy online visit [www.mtg.in](http://www.mtg.in).

For more information or for help in placing your order, call 0124-6601200 or e-mail: [info@mtg.in](mailto:info@mtg.in)



# ABRACADABRA



More than magic words, you need help from a magician to pass JEE with flying colours. Which is why MTG has collaborated with Dr Ramanalah Gundala, the popular Maths professor from Chennai, to bring you the all-new Objective Mathematics for JEE Main & Advanced.

## Includes

6 assignments for self-assessment  
Previous years' JEE questions (2017-13)

Introductory price  
₹ 750/-



Dr Ramanalah Gundala

After securing his DIIT and Ph.D from IIT Kharagpur, Dr Gundala was elected Fellow of National Academy of Sciences (FNASc). His 50+ years of teaching experience includes distinguished tenures at IIT Kharagpur and Anna University, Chennai. He has authored 7 books and published an astonishing 85 research papers. He's now retired and prepares students for success in IIT-JEE at two leading coaching institutes in Chennai and Warangal.

## Highlights of MTG's Objective Mathematics for JEE Main & Advanced

- Well-structured theory covering summary of important concepts and easy-to-understand illustrations
- 5800+ questions
- Unique and brain-stimulating exercises including questions of the following types:
  - Single Correct Option · More Than One Correct Option
  - Assertion and Reason. Comprehension Type
  - Matrix Match · Integer Answer Type

Visit [www.MTG.in](http://www.MTG.in) to buy online or visit a leading bookseller near you.  
For more information, e-mail [info@mtg.in](mailto:info@mtg.in) or call 1800 300 23355 (toll-free) today.

**mtg**

**TRUST OF MILLIONS, SINCE 1982**

# PRODUCING TOP RANKERS FOR THE LAST 29 YEARS

## WE BELIEVE YOU CAN

At Aakash, we have been nurturing students for last 29 years to help them excel in their career in Medical, Engineering & Foundations coaching. Enroll with us today to experience the best in class faculty and course material and build a successful career, because WE BELIEVE YOU CAN.

## ADMISSION OPEN

ONE YEAR / TWO YEAR INTEGRATED CLASSROOM COURSES 2019, 2020

### MEDICAL

**NEET & AIIMS /  
Other Medical Entrance Exams**

For Class XI students moving to Class XII and  
for Class X students moving to Class XI respectively

### ENGINEERING

**JEE (Main & Advanced) /  
Other Engg. Entrance Exams**

For Class XI students moving to Class XII and  
for Class X students moving to Class XI respectively

### FOUNDATIONS

**NTSE, Olympiads &  
School / Board Exams**

For Class IX & X students

### DIGITAL & DISTANCE LEARNING PROGRAMS



Printed Study Material



Live Online Classes



Recorded Video Lectures



# Aakash

**Medical | IIT-JEE | Foundations**

(Divisions of Aakash Educational Services Pvt. Ltd.)



[www.aakash.ac.in](http://www.aakash.ac.in)



TOLL FREE: **1800-313-0329**



HELPLINE NUMBER: **39454545**



GIVE A MISSED CALL: **9599698693**



SMS Aakash to **53030**

Registered Office : Aakash Tower, 8, Pusa Road, New Delhi-110005. Ph.: (011) 47623456 | E-mail: [iiitjee@aesl.in](mailto:iiitjee@aesl.in) | [medical@aesl.in](mailto:medical@aesl.in) | [info.afs@aesl.in](mailto:info.afs@aesl.in) | [dlp@aesl.in](mailto:dlp@aesl.in) | [aakashlive@aesl.in](mailto:aakashlive@aesl.in) | [aakashitutor@aesl.in](mailto:aakashitutor@aesl.in)



[instagram.com/aakasheducation](https://www.instagram.com/aakasheducation)



[facebook.com/aakasheducation](https://www.facebook.com/aakasheducation)



[youtube.com/AakashEducation](https://www.youtube.com/AakashEducation)



[twitter.com/aakash\\_twitted](https://twitter.com/aakash_twitted)



[aakashinstitute.blogspot.com](http://aakashinstitute.blogspot.com)